

BEET (SUGAR)

General Information

GENERAL

RANCONA Sugarbeets is a systemic broad-spectrum fungicide and seed dressing specifically for protection against soilborne and seedborne diseases. RANCONA Sugarbeets is active against Rhizoctonia and Fusarium. In addition, RANCONA Sugarbeets provides activity against fungi such as Mucor, Rhizopus, Aspergillus, Penicillium, Alternaria, Botrytis, and Cladosporium.

RANCONA Sugarbeets does not provide activity against Pythium spp. To provide best seedling protection use RANCONA Sugarbeets in combination with a product containing metalaxyl (such as Belmont 2.7 FS). Consult the label for appropriate use rate and follow all use instructions and precautions for both products. Do not use with other seed treatment products unless previous experience assures compatibility.

RANCONA Sugarbeets may be applied as a water-based slurry with other registered seed treatment insecticides and fungicides through standard slurry or mist-type commercial seed treatment equipment.

Not for use on agricultural establishments in hopper-box, planter-box, slurry-box or other seed treatment applications at or immediately before planting.

NOTE: This product is undyed. The purchaser of this product is responsible for ensuring that all seed treated with this product are adequately colored with a suitable colorant to prevent its accidental use as food for man or feed for animals. Refer to 21CFR, Part 2.25. Any colorant added to treated seed must be cleared for use under 40CFR, Part 153.155.

SEED TREATMENT EQUIPMENT

RANCONA Sugarbeets may be applied with mechanical, slurry, or mist-type seed treating equipment, provided that the equipment can be calibrated to accurately and uniformly apply the product to seed. Uniform application to seed is necessary to ensure best disease protection and optimum performance. Seed should be sound

and well-cured before treatment. Refer to the label rates below. Product requires dilution prior to application. Contact your local Chemtura representative or supplier for specific recommendations.

TREATED SEED LABELING

Seed that has been treated with this product that is then packaged or bagged for future use must contain the following labeling on the outside of the seed package or bag:

“This bag contains seed treated with ipconazole. Do not use for Food, Feed, or Oil Purposes. Persons opening this bag or loading/ pouring the treated seed must wear a long-sleeved shirt, long pants, shoes, socks and chemical-resistant gloves. After seeds have been planted, do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Exception: Once seeds are planted and covered by soil or other planting media, the Worker Protection Standard (WPS) allows workers to enter the treated area without restriction.” “Do not graze or feed livestock on treated areas”

FAILURE TO FOLLOW DIRECTIONS ON THIS LABEL MAY RESULT IN CROP INJURY OR POOR DISEASE CONTROL.

Method

[Seed Treatment](#)

Rates

[field rates 0](#)

•

Restricted Entry Interval

12 hours

Exception: Once seeds are planted and covered by soil or other planting media, the Worker Protection Standard (WPS) allows workers to enter the treated area without restriction.”

Timings

[N.A.](#)