

# **TURF**

## General Information

### GENERAL INFORMATION

VITICULTURE ZINC can be used in combination with water-soluble plant nutritional products which are formulated for foliar, sidedress or drip irrigation application. May be applied alone or in tank mixes containing fertilizers, fungicides and/or pesticides. VITICULTURE ZINC is believed to be compatible with most agricultural water-soluble plant

nutritional products. If physical compatibility with another fertilizer product is unknown, use

a jar compatibility test.

The addition of calcium to the VITICULTURE ZINC is recommended.

When applying VITICULTURE ZINC blend as a foliar spray, always use a spreader penetrant.

ONCE PRODUCT IS DILUTED, USE WITHIN 24 HOURS

### DIRECTIONS FOR USE

Apply as needed based on soil tests or according to recommendations in your nutrient

management plan. Also refer to Suggested Use Rates below.

Manufacturer Use Product: This product is to be used only in combination with other fertilizers.

Citrus & Avocados: 1.0 to 4.0 quarts per acre. Apply after first flush in Spring. Repeat applications every 2 to 4 weeks as necessary.

Deciduous Fruit & Nut Crops: 1.0 to 4.0 quarts per acre. Apply as needed throughout the growth season.

Field & Vegetable Crops: 1.0 to 4.0 quarts per acre. Begin applications when plants are

6 to 8 inches in size.

Grapes & Berries: 1.0 to 4.0 quarts per acre. Apply as needed during periods of active

growth.

Ornamental Tree, shrubs and flowering plants: Apply at dilution rate of 3.0 ounces in 1.0 gallon of water. Spray to runoff. Apply as needed at spray intervals of 14 days or

more.

Rates recommended are for standard spray volumes of 50.0 to 150 gallons per acre (500

to 1,500 liters per hectare). For higher volume sprays, rate should be adjusted.

Turf: 0.5 to 1.0 fluid ounces per 1,000 sq. ft. Weekly applications as needed.

Caution: Store Viticulture Zinc in temperatures between 41 °F to 104 °F and avoid extreme variations in temperature. A reversible separation of ingredients may occur after

long storage, this however does not affect quality or effectiveness of the product.

Limitations, Restrictions, and Exceptions

Turf: 0.5 to 1.0 fluid ounces per 1,000 sq. ft. Weekly applications as needed.

Rates

[field\\_rates 0](#)

- 

Timings

[Weekly applications as needed](#)