

FOLIAR

General Information

For use as a fertilizer in foliar and fertigation applications to supplement nutrient levels on agricultural crops.

DIRECTIONS FOR USE (Use only as Directed):

- Exercise caution with foliar applications in extreme weather conditions such as high temperatures and poor drying conditions.
- Aerial application - minimum of 5 gallons of water/acre; Dilute sprays - minimum of 20 gallons of water/acre. Apply with sufficient water to deliver spray uniformly on leaf surfaces. Higher volumes of water should be used for higher rates of fertilizer.
- Exercise caution when applying to any crop during bloom, as many factors can affect fruit set and final fruit quality.
- BudMax is compatible with most agricultural chemicals and nutrients. Not compatible with aluminum equipment. Should be applied in conjunction with tissue and/or soil analysis.
- Application rates vary according to specific crop requirements, management objectives, soil nutrient availability and/or environmental conditions.
- Consult a NutriAg representative for specific crop recommendations, application timings and necessity of its use.

ATTENTION: This fertilizer contains boron (B). The use of this fertilizing material on any crops other than those recommended may result in serious injury to crop(s).

ATTENTION: This fertilizer contains molybdenum (Mo). The application of fertilizing materials containing molybdenum (Mo) may result in forage crops containing levels of molybdenum (Mo) which are toxic to ruminant animals.

Method

[Broadcast/Foliar Air](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

-

Timings

[At pink bud to early bloom, post-nut set, and/ or late season during following year bud differentiation.](#)