

OUTDOOR PERIMETER BARRIER TREATMENT AROUND STRUCTURES - 2 PACKET

General Information

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Exit area immediately and remain outside the treated area until sprays have dried.

IMPORTANT: Not for use on plants being grown for sale or other commercial use, or for commercial seed production or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks or on lawns or grounds.

For best results, thoroughly wash out sprayer and screen with water and detergent before using Cyper WSP.

INDOOR USE RESTRICTIONS

- Do not use water-based sprays in or on conduits, motor housings, junction boxes, or other electrical equipment because of possible shock hazard.
- During any indoor surface application, do not allow dripping or run-off to occur. During any application to ceilings of a structure, cover surface below with plastic shielding or similar material.
- Do not apply this product in any room being used as a living, eating, or recovery area by patients, the elderly, or the infirm when they are in the room.
- Do not apply to classrooms when in use.
- Do not apply to areas of institutions (including libraries sports facilities, etc.) when occupants are present in the immediate treatment area.
- Do not use as a space spray. Use only in well-ventilated areas.
- Do not use concentrate or emulsion in fogging equipment.
- Do not use in food areas of food-handling establishments, which includes areas for receiving, serving storage (dry, cold, frozen raw), packaging (canning, bottling, wrapping, boxing), preparing (cleaning, slicing, cooking, grinding), edible waste

storage, enclosed processing systems (mills, dairies, edible oils, syrups). In the home, all food-processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed.

- Do not use in warehouses while raw agricultural commodities for food or feed and/or raw or cured tobacco are being stored. Do not use in greenhouses where crops for food or feed are grown.

Pet Restrictions

- Do not apply to pets. Remove birds and other pets before application. Cover any water inhabited by fish (such as aquariums and ornamental fish ponds) during treatment and turn aquarium systems off.

OUTDOOR USE RESTRICTIONS

- All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses.

1. Treatment to soil or vegetation around structures,
2. Applications to lawns, turf, and other vegetation
3. Applications to building foundations up to a maximum height of 3 feet.

- All outdoor applications to impervious surfaces such as sidewalks, driveways, patios porches, and structural surfaces (such as windows, doors, and eaves) are limited to spot treatments or crack-and-crevice applications only.

- Do not apply directly to sewers or drains or to any area like a gutter where drainage to sewers storm drains, water bodies or aquatic habitat can occur.

Applications around Swimming Pools

- Do not apply directly to swimming pools or swimming pool systems.

- This product may be applied as a broadcast treatment to lawns and other vegetated areas around swimming pools, or as a spot treatment or crack-and-crevice treatment to impermeable surfaces (such as tiled walkways) around pools

Application in and on Boats

- Do not apply to boat surfaces which contact water. Broadcast applications to exterior surfaces of boats are prohibited Spot treatments may be made to exterior surfaces that do not contact water.

- Use inside boats, ships, and other vessels is permitted. Do not allow product to drain or wash off into water bodies or other aquatic habitat.

Applications near or around Aircraft

- Do not apply to aircraft cabins

USE INFORMATION

For use as a general surface crack-and-crevice and/or spot treatment for residual and contact pest control in on and around buildings and structures and their immediate surroundings and on modes of transport. Permitted areas of use in nonfood/nonfeed areas include but are not limited to, aircraft (cargo and non-cabin areas only), apartment buildings boiler rooms, buses, closets, correctional facilities decks entries, factories fencing, floor drains (that lead to sewers), food granaries, food grain mills, food manufacturing, processing and serving establishments, furniture, garages, garbage rooms greenhouses (non-commercial) hospitals (non-patient areas) hotels and motels houses industrial buildings laboratories landscape vegetation, locker rooms machine rooms, mausoleums, mobile homes, mop closets mulch, nursing homes, offices patios pet kennels, porches, railcars, restaurants, storage rooms schools sewers (dry) stores trailers, trees, trucks, utility passages, vessels, vestibules warehouses, wineries and yards.

Food-Handling Establishments (For Use In Non-Food Areas Only):

Cyper WSP may be used as a general spot or crack and crevice treatment in nonfood areas of food handling establishments. Examples of nonfood areas include garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets, and storage (after canning or bottling).

Residential Food Areas

In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed.

Application Equipment Preparation

For best results thoroughly wash out sprayer and screens with water and detergent before using Cyper WSP. To avoid sprayer clogging do not use strainers or screens finer than 50 mesh size.

INDOOR SURFACES USE

Cyper WSP may be used as a crack-and-crevice and/or spot treatment for indoor surfaces.

OUTDOOR SURFACE USE

Apply with either hand-held or power application equipment as a residual spray to surfaces of buildings, porches, screens, window frames, eaves, patios, lawns, refuse dumps, garages and other similar areas where these pests are found. Also, may be used as a spot treatment for applications to lawns in landscaped areas in and around residential, commercial, institutional and industrial buildings, and similar areas where these insect pests are active.

Note: The maximum application rate to lawns and other turfgrass is 0.44 lb. ai/A (0.0101 lb. ai/1,000 sq. ft.).

Cluster Flies

Make exterior treatments in late summer or early fall before flies are observed alighting on surfaces. Apply thoroughly on siding, under eaves, and around windows and doors paying particular attention to south-facing surfaces. Apply just enough dilution to adequately cover the area without excessive dripping or runoff. Heavy precipitation prior to frost may require retreatment to maintain protection.

Limitations, Restrictions, and Exceptions

OUTDOOR PERIMETER BARRIER TREATMENT AROUND STRUCTURES

Applying a continual band of insecticide around building foundation and around windows, doors, service lines, entrances, eaves, vents, and other areas can greatly reduce the potential for entry by crawling pests. To facilitate application, remove debris and leaf litter from next to the foundation, trim vegetation and branches that touch the foundation and move or rake back rocks, deep mulch or other potential pest harborage next to the foundation. To help prevent infestation of buildings, apply as band to soil and vegetation 6-10 ft. wide around and adjacent to the building. Also, treat the building foundation to a height of 2-3 ft. where pests are active and may find entrance. Apply as a coarse spray treatment to thoroughly and uniformly wet the foundation and/or band area, using 1-5 gals. of treatment solution applied to 800-1600 sq. ft.

Amount of concentration is dependent upon species (see pest table and comments), infestation levels and service interval desired.

- Packet = 0.33 oz. (9.5 gms) of Cyper WSP (equivalent to 2 level TB)
- Application volume may be greater than 5 gal/800-1,600 sq. ft. if required under

heavy vegetative or landscaping materials in order to obtain desired coverage.

Ants

Apply to any trails, around doors and windows and similar areas where ants (including carpenter ants) may be found. Where ants are trailing inside, apply as a residual surface treatment to active areas such as baseboards, corners around pipes, in and behind cabinets, behind and under refrigerators, sinks, furnaces and stoves, cracks and crevices. When combining baits and residual surface insecticides, apply surface insecticides in cracks and crevices, along baseboards, and infested surfaces and outside barrier treatments. Apply perimeter barrier treatments and treat nest to prevent infestation as described below in Outdoor Surface Use.

Carpenter Ants

Apply as a coarse wet spray treatment, but avoid runoff around doors, frames, windows and similar areas where these ants enter premises or hide.

Bees, Flies, Wasps

Apply directly to walls, window screens and other exterior resting areas as a residual surface treatment. Use caution when treating nests of stinging insects as Cyper WSP does not provide instant knockdown, Protective equipment for the applicator may be required. For best results, treat bee, wasp and hornet nests late in the day when most insects will be present.

Allow 2-3 days for the colony to die.

Cockroaches, Spiders, Ticks

Apply as a coarse, low pressure treatment to areas where these pests hide, such as baseboards, corners, storage areas, closets, around water pipes, doors and windows, attics and eaves, cabinets, sinks, furnaces, stoves, behind and under refrigerators, the underside of shelves, drawers and similar areas. Pay particular attention to cracks and crevices; also see Outdoor Surface Use.

Ticks

To control nuisance fleas and ticks (e.g. dog ticks) apply to yards, runs and other outside areas where pets may frequent. To control ticks, apply using a coarse fan spray to vegetation, brush, branches, rock walls, and other areas near habitation where ticks may harbor or frequent. Treat entire area rather than making spot treatments, and retreat as necessary to maintain control.

Do not apply to pasture or cropland, and do not allow animals and people access to treated areas until the sprays have dried. Applications can begin in the spring and can continue until frost to control both larvae and adult ticks.

Scorpions

Treat and remove accumulations of lumber, firewood and other materials which serve as harborage sites. Apply a coarse spray treatment to the point of drip.

Method

[Band](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

-

Timings

[N.A.](#)