

TERMITES CONTROL (NOT FOR SOIL TREATMENT) - TERMITES CARTON NEST IN TRESS OR BUILDING VOIDS

General Information

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Exit area immediately and remain outside the treated area until sprays have dried.

IMPORTANT: Not for use on plants being grown for sale or other commercial use, or for commercial seed production or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks or on lawns or grounds.

For best results, thoroughly wash out sprayer and screen with water and detergent before using Cyper WSP.

INDOOR USE RESTRICTIONS

- Do not use water-based sprays in or on conduits, motor housings, junction boxes, or other electrical equipment because of possible shock hazard.
- During any indoor surface application, do not allow dripping or run-off to occur. During any application to ceilings of a structure, cover surface below with plastic shielding or similar material.
- Do not apply this product in any room being used as a living, eating, or recovery area by patients, the elderly, or the infirm when they are in the room.
- Do not apply to classrooms when in use.
- Do not apply to areas of institutions (including libraries sports facilities, etc.) when occupants are present in the immediate treatment area.
- Do not use as a space spray. Use only in well-ventilated areas.
- Do not use concentrate or emulsion in fogging equipment.
- Do not use in food areas of food-handling establishments, which includes areas for receiving, serving storage (dry, cold, frozen raw), packaging (canning, bottling, wrapping, boxing), preparing (cleaning, slicing, cooking, grinding), edible waste

storage, enclosed processing systems (mills, dairies, edible oils, syrups). In the home, all food-processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed.

- Do not use in warehouses while raw agricultural commodities for food or feed and/or raw or cured tobacco are being stored. Do not use in greenhouses where crops for food or feed are grown.

Pet Restrictions

- Do not apply to pets. Remove birds and other pets before application. Cover any water inhabited by fish (such as aquariums and ornamental fish ponds) during treatment and turn aquarium systems off.

OUTDOOR USE RESTRICTIONS

- All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses.

1. Treatment to soil or vegetation around structures,
2. Applications to lawns, turf, and other vegetation
3. Applications to building foundations up to a maximum height of 3 feet.

- All outdoor applications to impervious surfaces such as sidewalks, driveways, patios porches, and structural surfaces (such as windows, doors, and eaves) are limited to spot treatments or crack-and-crevice applications only.

- Do not apply directly to sewers or drains or to any area like a gutter where drainage to sewers storm drains, water bodies or aquatic habitat can occur.

Applications around Swimming Pools

- Do not apply directly to swimming pools or swimming pool systems.

- This product may be applied as a broadcast treatment to lawns and other vegetated areas around swimming pools, or as a spot treatment or crack-and-crevice treatment to impermeable surfaces (such as tiled walkways) around pools

Application in and on Boats

- Do not apply to boat surfaces which contact water. Broadcast applications to exterior surfaces of boats are prohibited Spot treatments may be made to exterior surfaces that do not contact water.

- Use inside boats, ships, and other vessels is permitted. Do not allow product to drain or wash off into water bodies or other aquatic habitat.

Applications near or around Aircraft

- Do not apply to aircraft cabins

USE INFORMATION

For use as a general surface crack-and-crevice and/or spot treatment for residual and contact pest control in on and around buildings and structures and their immediate surroundings and on modes of transport. Permitted areas of use in nonfood/nonfeed areas include but are not limited to, aircraft (cargo and non-cabin areas only), apartment buildings boiler rooms, buses, closets, correctional facilities decks entries, factories fencing, floor drains (that lead to sewers), food granaries, food grain mills, food manufacturing, processing and serving establishments, furniture, garages, garbage rooms greenhouses (non-commercial) hospitals (non-patient areas) hotels and motels houses industrial buildings laboratories landscape vegetation, locker rooms machine rooms, mausoleums, mobile homes, mop closets mulch, nursing homes, offices patios pet kennels, porches, railcars, restaurants, storage rooms schools sewers (dry) stores trailers, trees, trucks, utility passages, vessels, vestibules warehouses, wineries and yards.

Food-Handling Establishments (For Use In Non-Food Areas Only):

Cyper WSP may be used as a general spot or crack and crevice treatment in nonfood areas of food handling establishments. Examples of nonfood areas include garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets, and storage (after canning or bottling).

Residential Food Areas

In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed.

Application Equipment Preparation

For best results thoroughly wash out sprayer and screens with water and detergent before using Cyper WSP. To avoid sprayer clogging do not use strainers or screens finer than 50 mesh size.

INDOOR SURFACES USE

Cyper WSP may be used as a crack-and-crevice and/or spot treatment for indoor surfaces.

OUTDOOR SURFACE USE

Apply with either hand-held or power application equipment as a residual spray to surfaces of buildings, porches, screens, window frames, eaves, patios, lawns, refuse dumps, garages and other similar areas where these pests are found. Also, may be used as a spot treatment for applications to lawns in landscaped areas in and around residential, commercial, institutional and industrial buildings, and similar areas where these insect pests are active.

Note: The maximum application rate to lawns and other turfgrass is 0.44 lb. ai/A (0.0101 lb. ai/1,000 sq. ft.).

Cluster Flies

Make exterior treatments in late summer or early fall before flies are observed alighting on surfaces. Apply thoroughly on siding, under eaves, and around windows and doors paying particular attention to south-facing surfaces. Apply just enough dilution to adequately cover the area without excessive dripping or runoff. Heavy precipitation prior to frost may require retreatment to maintain protection.

Limitations, Restrictions, and Exceptions

Termites Control (Not for Soil Treatment)

For control of termites, subterranean aerial colonies, Formosan aerial colonies or drywood termites in localized areas of infested wood structures, apply to voids and galleries in damaged wood and in spaces between wooden members of a structure and between wood and foundations where wood is vulnerable. Application may be made to inaccessible areas by drilling, and then injecting the emulsion with a crack and crevice injector into the damaged wood or void spaces. This type of application is not intended to be a substitute for soil treatment for extensive infestation of drywood termites or other wood-infesting insects. This treatment is intended to kill localized infestations of workers and winged reproductive forms of termites.

Termite carton nests in trees or building voids may be injected with 0.25%-0.50% solution using a pointed injection tool. Multiple injection points to varying depths may be necessary. It is desirable to physically remove carton nest material from building voids when such nests are found.

All leaks resulting from the deposition of termiticide in locations other than those prescribed on this label must be cleaned up prior to leaving the application site.

Use anti-backflow equipment or procedures to prevent siphonage of pesticide back

into water supplies.

Care should be taken that the treatment solution is not introduced into the gravel and/or pipe drainage system which may be located on the exterior of the foundation in close proximity to the footing of the structure.

RESTRICTIONS:

- Do not use this product for pre-construction soil treatment under concrete slabs.
- Do not allow people or pets to contact contaminated areas or to reoccupy the contaminated area until the clean-up is completed.
- Do not treat soil that is water-saturated or frozen.
- Do not treat when raining.
- Do not allow treatment to run off from the target area.
- Do not apply within 10 feet of storm drains.
- Do not apply within 25 feet of aquatic habitats (such as, but not limited to lakes, reservoirs, rivers, permanent streams, marshes, or natural ponds, estuaries, and commercial fish farm ponds).
- Do not make on-grade applications when sustained wind speeds are above 10 mph (at application site) at nozzle end height.

Method

[Injection](#)

Rates

[field rates 0](#)

•

Timings

[N.A.](#)