

INDOOR TREATMENTS - DRAIN FLIES

General Information

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

USE RESTRICTIONS

- Apply this product only as specified on this label.
- DO NOT compost any discarded plant materials that have been treated with this product.
- DO NOT make outdoor applications during rain.
- DO NOT reapply to treated areas more than 1x every 4 weeks.
- DO NOT allow adults, children or pets to enter until vapors, mists and/or aerosols have dispersed and the treated area has been thoroughly ventilated. DO NOT contact treated surfaces until dry.
- When used in schools, DO NOT apply to classrooms while in use. Allow treated areas to dry thoroughly before reoccupying.
- When used in hospitals and nursing homes, remove patients before treating rooms. Allow treated areas to dry thoroughly before returning patients to rooms.
- When used in the home, remove or cover exposed food and drinking water in the treatment area before application. Remove or cover dishes, utensils, food processing equipment and food preparation surfaces in the treatment area or wash them before use. DO NOT use in serving areas while food is exposed.
- Applications in food areas of food handling establishments are limited to spot and/or crack and crevice treatments. DO NOT make applications when facility is in operation. Avoid contamination of food or feedstuffs. DO NOT apply directly to food, food packaging or food contact surfaces. During application, cover or remove food

and cover food processing surfaces and equipment, or clean after treatment with a suitable detergent and rinse with potable water before use. DO NOT use in serving areas (dining areas where prepared foods are served) while food is exposed or facility is in operation.

- Except when applying to sewers or drains as instructed on this label, DO NOT apply directly into sewers or drains or to any area like a gutter where drainage to storm sewers, water bodies, or aquatic habitat can occur.
- When used in zoos, remove animals and cover or remove animal food/water prior to application.
- Remove pets and cover or remove pet food/water and fish bowls prior to application.

CRACK & CREVICE TREATMENT: Use with the supplied actuator and injection tubes. Inject into cracks and crevices or void spaces where insects may be harboring, living and breeding. Place injector tip into cracks, crevices, holes and other small openings. Release approximately 1 sec of product. For light infestations, move injector tip along cracks while treating at the rate of 3 In ft/sec. For heavy infestations, move injector tip along at 1 In ft/sec. For closed voids, calculate the void's cubic area and treat at the rate of 1 to 5 sec/3 ft³. Several holes may be required in long-running voids.

SPOT TREATMENT: Use with the supplied actuator and injection tubes. Hold container upright while spraying. Direct nozzle approximately 12 inches from surface to be treated. Apply directly on insects in these locations when possible. Limit individual spot treatments to an area no larger than 2 ft²/room.

Before spot treating on plastic, painted or varnished surfaces, test a small, inconspicuous area and observe for staining or damage to finish.

Limitations, Restrictions, and Exceptions

DRAIN FLIES: Release approximately 1 to 2 sec of product into the drain basin.

Method

[Injection](#)

Timings

N.A.