

# **LIVESTOCK HOUSING - DARKLING BEETLES**

## General Information

### DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

### USE RESTRICTIONS

- DO NOT apply to humans, their clothing or bedding.
- Cover fish tanks prior to application.
- DO NOT contaminate food, feedstuffs, or water supply. DO NOT contaminate food preparation surfaces, kitchen utensils, dishes or feed storage containers. Cover all food contact surfaces and cooking utensils during treatment or thoroughly wash with an effective cleaning compound followed by a potable water rinse.

### INFORMATION

This product is a flowable microencapsulated concentrate containing chlorpyrifos. It may be sprayed on any surface which will not be damaged or stained by water. Heavy applications may leave a visible deposit. On some surfaces this deposit can be removed with a damp cloth or sponge.

Mixing Instructions: Thoroughly clean spray equipment before using this product. Shake concentrate well before diluting. Apply at the rate of 2 to 5 fl oz of concentrate per 1,000 ft<sup>2</sup>, using an adequate amount of water to ensure good coverage. Flies require a higher application rate. See the “Flies” section below for specific directions. When diluting, first add approximately 1/2 of the water to the spray tank and then add the proper amount of this product. Add the rest of the water and agitate the sprayer thoroughly. Agitate sprayer occasionally during use to ensure even coverage. Shake or reagitater sprayer if spraying is interrupted or if dilution is left in sprayer overnight. When emptying the bottle, triple rinse with water and add the rinsate to the spray tank. If spray screens are used, they should be 50 mesh or larger. In situations where conventional application methods have not or are not likely to provide adequate coverage, foam generating equipment or similar machines can be used to provide a continuous coverage. Use drier foam inside wall voids to avoid any water damage or electrical short circuits.

#### Limitations, Restrictions, and Exceptions

#### LIVESTOCK HOUSING

FOR USE: Livestock Housing and Holding Areas (such as Hog Barns, Empty Chicken Houses, Dairy Areas, Milkrooms, Calf Hutches, Calving Pens and Parlors).

DO NOT apply to lactating or food animals. DO NOT contaminate milk, food or drinking water. Remove all animals prior to making application to housing. DO NOT apply or allow insecticide to drift onto animals. DO NOT allow animals to enter treated areas until surfaces are dry.

DARKLING BEETLES: Treat the litter paying particular attention to the areas under the feeder line, water line and the perimeter. Use 2 to 5 fl oz/1,000 ft<sup>2</sup> in an adequate amount of water to provide good coverage.

Method

[N.A.](#)

Timings

[N.A.](#)