

# **COMMERCIAL INDOOR TREATMENTS - CARPET BEETLES (DERMESTIDS)**

## General Information

### DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

### USE RESTRICTIONS

- DO NOT apply to humans, their clothing or bedding.
- Cover fish tanks prior to application.
- DO NOT contaminate food, feedstuffs, or water supply. DO NOT contaminate food preparation surfaces, kitchen utensils, dishes or feed storage containers. Cover all food contact surfaces and cooking utensils during treatment or thoroughly wash with an effective cleaning compound followed by a potable water rinse.

### INFORMATION

This product is a flowable microencapsulated concentrate containing chlorpyrifos. It may be sprayed on any surface which will not be damaged or stained by water. Heavy applications may leave a visible deposit. On some surfaces this deposit can be removed with a damp cloth or sponge.

Mixing Instructions: Thoroughly clean spray equipment before using this product. Shake concentrate well before diluting. Apply at the rate of 2 to 5 fl oz of concentrate per 1,000 ft<sup>2</sup>, using an adequate amount of water to ensure good coverage. Flies require a higher application rate. See the "Flies" section below for specific directions. When diluting, first add approximately 1/2 of the water to the spray tank and then add the proper amount of this product. Add the rest of the water and agitate the sprayer thoroughly. Agitate sprayer occasionally during use to ensure even coverage. Shake or reagitate sprayer if spraying is interrupted or if dilution is left in sprayer overnight. When emptying the bottle, triple rinse with water and add the rinsate to the spray tank. If spray screens are used, they should be 50 mesh or larger. In situations where conventional application methods have not or are not likely to provide adequate coverage, foam generating equipment or

similar machines can be used to provide a continuous coverage. Use drier foam inside wall voids to avoid any water damage or electrical short circuits.

## Limitations, Restrictions, and Exceptions

### COMMERCIAL INDOOR TREATMENTS

FOR USE IN: Non-Food Areas of Manufacturing, Industrial Plants; Warehouses; Ship Holds; Railroad Boxcars. Apply at the rate of 2 to 5 fl oz of concentrate per 1,000 ft<sup>2</sup>, using an adequate amount of water to ensure good coverage. DO NOT use in food areas of food processing plants.

SAMPLE SPRAY DILUTION CHART: Examples of possible dilutions using this product with water are given in the table at the label.

A higher rate should be used for clean-outs or for high insect infestations.

Use as a Crack & Crevice product where insects may be hiding, living and breeding and/or for selected spot treatment.

CARPET BEETLES (DERMESTIDS): Thoroughly apply the spray as a spot treatment along edges of carpet, under carpet edge (if necessary), under furniture and throw rugs. Use as a Crack & Crevice application in closets and other shelving where infested items are stored. DO NOT treat clothing with this product. Repeat application as necessary.

Method

[Spot treatment](#)

Timings

[N.A.](#)