

COMMERCIAL OUTDOOR BUILDING TREATMENTS - OUTDOOR COMMERCIAL USE

General Information

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

USE RESTRICTIONS

- DO NOT apply to humans, their clothing or bedding.
- Cover fish tanks prior to application.
- DO NOT contaminate food, feedstuffs, or water supply. DO NOT contaminate food preparation surfaces, kitchen utensils, dishes or feed storage containers. Cover all food contact surfaces and cooking utensils during treatment or thoroughly wash with an effective cleaning compound followed by a potable water rinse.

INFORMATION

This product is a flowable microencapsulated concentrate containing chlorpyrifos. It may be sprayed on any surface which will not be damaged or stained by water. Heavy applications may leave a visible deposit. On some surfaces this deposit can be removed with a damp cloth or sponge.

Mixing Instructions: Thoroughly clean spray equipment before using this product. Shake concentrate well before diluting. Apply at the rate of 2 to 5 fl oz of concentrate per 1,000 ft², using an adequate amount of water to ensure good coverage. Flies require a higher application rate. See the "Flies" section below for specific directions. When diluting, first add approximately 1/2 of the water to the spray tank and then add the proper amount of this product. Add the rest of the water and agitate the sprayer thoroughly. Agitate sprayer occasionally during use to ensure even coverage. Shake or reagitate sprayer if spraying is interrupted or if dilution is left in sprayer overnight. When emptying the bottle, triple rinse with water and add the rinsate to the spray tank. If spray screens are used, they should be 50 mesh or larger. In situations where conventional application methods have not or are not likely to provide adequate coverage, foam generating equipment or

similar machines can be used to provide a continuous coverage. Use drier foam inside wall voids to avoid any water damage or electrical short circuits.

Limitations, Restrictions, and Exceptions

COMMERCIAL OUTDOOR BUILDING TREATMENTS

OUTDOOR COMMERCIAL USE: To help prevent infestations of buildings, treat a band of soil 6 to 10 ft wide around and adjacent to the building. Also treat the building foundation to a height of 2 to 3 ft and where pests are active and may find entrance. Treat other insect entry points around doors, windows, under siding and eaves of structure, under decks and on fences. This product may be applied in compressed air equipment or power spray equipment. Depending on type of equipment used to apply product and landscaping of perimeter, use 0.5 to 1.75 fl oz/1,000 ft². Alternate directions are to use 5.0 to 8.75 fl oz of product per 50 gal of water applied at 10 gal of dilution per 1,000 ft².

THIS PRODUCT DOES NOT PROVIDE INSTANT KNOCKDOWN. THE BEST TIME TO TREAT IS LATE IN THE DAY. COLONY SHOULD BE DEAD IN 2 TO 3 DAYS. PROTECTIVE EQUIPMENT FOR THE APPLICATOR IS REQUIRED.

Method

[Spray](#)

Timings

[N.A.](#)