

LATE SEASON PLANT AND SOIL SURFACE TREATMENTS

General Information

High Concentrate Dark Turf Colorant For Seasonal Growth Transitions

- Highest strength formulation for application flexibility
- Extends late-fall and early-winter transitional play
- Apply to warm and cool season turfgrasses in all regions

Green Lawngr Transition HC is a high concentrate dark turf colorant made with ColorLock for applications on all types of turfgrasses in all growing regions.

Transition HC is for the turfgrass professional who wants to enhance the aesthetics of pre-dormant and dormant warm- and cool-season turf varieties. Unlike dyes and other colorants that fade quickly or turn from green to blue, Transition HC dark turf colorant produces the most uniform and consistent long lasting color with dramatic effects. The high concentrate formulation of Transition HC gives the turfgrass professional greater color range and application flexibility to achieve just the right custom look.

Directions For Use

Recommended Rates: For most dormant turfgrass varieties, begin applications at 32 ounces per acre. Pre-dormant (late fall to early winter) applications may start at a higher rate of 64 ounces per acre to create a base color on existing turf. As frost events occur and dormancy progresses, maintaining or gradually reducing the applications rates of Transition HC will create a more natural look of turf transition. One (1) gallon of Transition HC will adequately treat up to 4 acres of close-mowed dormant turf. For the most uniform coverage, apply Transition HC with flat fan spray nozzles in a spray volume of 1-2 gallons of mix solution per 1,000 square feet of area (45-90 gallons per acre).

Limitations, Restrictions, and Exceptions

Late Season Plant and Soil Surface Treatments

Apply Transition HC at a rate of 16-64 ounces per acre from pre-dormancy to dormancy. When applied at the higher rate prior to dormancy, Transition HC will act to absorb UV light energy and help elevate plant canopy and soil surface temperatures. Transition HC may be applied every two to four weeks through the

dormancy period to maintain the desired turf color.

Method

[Soil Surface Treatment](#)

Timings

[From pre-dormancy to dormancy.](#)