

ALFALFA (GROWN FOR SEED) - SUPPLEMENTAL LABEL

General Information

DIRECTIONS FOR USE

Resistance:

Any insect/mite population may contain individual insects that may develop resistance to a specific pesticide product used in consecutive generations to control these pests. Prediction of resistance development is uncertain. Follow appropriate resistance management strategies. Contact your local extension specialist or certified crop advisors for any additional pesticide resistance management and/or Integrated Pest Management recommendations for the specific site and pest problems in your area.

If resistance to this product develops in your area, you may find reduced control from this product or other products with a similar mode of action. If poor performance cannot be related to improper application methods or extreme weather, it is possible that a resistant strain of insect may be present. If poor control occurs and resistance is a reasonable cause, immediately consult with your local UPI representative or agricultural advisor for the best alternative method of control for your area.

Chemigation Use Directions

Apply this product only through sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation system. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system.

Use a minimum of 0.75 inch of water per acre for LEPA irrigation. Use 1 to 2 pints per acre of non-emulsified oils when used as the diluent. The set up and calibration of chemigation equipment are important to achieve control of target insect pests. Crop injury, lack of effectiveness, or illegal residues in the crop can result from non-uniform distribution of treated water. Contact your State Agricultural Extension Service specialists, equipment manufacturers or other experts for advice on the suitability of the equipment set up for optimum control of the target insect pests.

A person knowledgeable of the chemigation system and responsible for its operations, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise. Failure to cease application during a mechanical stoppage may result in undesirable residues to adjacent areas.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Apply Bifenture EC Agricultural Insecticide continuously for the duration of the water application. Dilute Bifenture EC Agricultural Insecticide in sufficient volume to ensure accurate application over the area to be treated. Use a minimum of 0.5 inch per acre of irrigation water for application by chemigation. No agitation is required if a suitable diluent is used. Conduct a diluent test to check that phase separation will not occur during dilution and application. If uniform dilution is not achieved during the entire period of application, undesirable residues or less than desirable control may occur.

Rotation Crop Restrictions

Crops for which bifenthrin tolerances exist may be rotated at any time. All other crops may be rotated 30 days after the last application.

Instructions for Application

The following Directions for Use tables provide, in some cases, rate ranges. Use the lower application rates when insect pressure is light to moderate. Use the higher application rates when insect populations are high, when treating mites and when climate is arid.

Cultivation within 10 feet of a water body is prohibited to allow for the growth of a vegetated filter strip.

In New York State this product may not be applied within 100 feet (using ground equipment) to 300 feet (using aerial equipment) of coastal marshes or permanent streams that drain into coastal marshes.

Limitations, Restrictions, and Exceptions

ALFALFA (Grown for Seed)

APPLICATION INSTRUCTIONS

- Application by ground: Apply specified dosage in a minimum of 20 gallons of finished spray per acre.
- Application by air: Apply specified dosage in a minimum of 5 gallons per acre. When applied by air, 1- 2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray.

Restrictions and Precautions

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product to blooming crops or weeds if bees are actively visiting the treatment area.

This pesticide is extremely toxic to fish and aquatic invertebrates. Use with care when applying in areas adjacent to any body of water. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark.

Do not make applications when weather conditions favor drift from treated areas. Drift and run-off from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

Apply when the wind velocity favors on target deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph.

Do not apply more than 6.4 ounces (0.1 lb active ingredient) per acre per application. Do not make more than 3 applications in a crop year.

Do not apply more than 19.2 ounces (0.3 lb. active ingredient) per acre per season.

Maintain a minimum of 21 days between applications. Rotational crops may be planted no sooner than 30 days after application.

This product may be applied through irrigation systems. Refer to the EPA registered label for chemigation directions.

Buffer Zones

Vegetative Buffer Strip: Construct and maintain a minimum 10-foot wide vegetative filter strip of grass or other permanent vegetation between the field edge and down gradient aquatic habitat (such as, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes or natural ponds; estuaries; and commercial fish farm ponds).

Only apply products containing bifenthrin onto fields where a maintained vegetative buffer strip of at least 10 feet exists between the field and down gradient aquatic habitat.

For guidance, refer to the following publication for information on constructing and maintaining effective buffers: Conservation Buffers to Reduce Pesticide Losses. Natural Resources Conservation Services. USDA, NRCS. 2000. Fort Worth, Texas. 21 pp. <http://www.in.nrcs.usda.gov/technical/agronomy/newconbuf.pdf>.

Buffer Zone for Ground Application (groundboom, overhead chemigation, or airblast): Do not apply within 25 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, permanent streams, marshes, natural ponds, estuaries, and commercial fish ponds).

Buffer Zone for ULV Aerial Application: Do not apply within 450 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, permanent streams, marshes, natural ponds, estuaries, and commercial fish ponds).

Buffer Zone for Non-ULV Aerial Application: Do not apply within 150 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, permanent streams, marshes, natural ponds, estuaries, and commercial fish ponds).

Additional Requirements for Ground Applications

- Wind speed must be measured adjacent to the application site on the upwind side, immediately prior to applications.
- For ground boom applications, apply using a nozzle height of no more than 4 feet above the ground or crop canopy.

Additional Requirements for Aerial Applications

- The spray boom should be mounted on the aircraft as to minimize drift caused by wingtip or rotor vortices. The minimum practical boom length should be used and must not exceed 75% of the wing span or 80% rotor diameter.
- Flight speed and nozzle orientation must be considered in determining droplet size.
- Spray must be released at the lowest height consistent with pest control and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety.
- When applications are made with a cross-wind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind

edge of the application area by adjusting the path of the aircraft upwind

Note: For use only on fields in production of alfalfa seed. Not for use on fields producing alfalfa for livestock. Seed from any treated crop on this label may not be used for human or animal consumptions. Alfalfa seed from treated fields may not be used for growing sprouts for humans or animal consumption. Producers of alfalfa grown for seed that use this product, or cause the product to be used on fields they operate, are required to inform, in writing, conditioners receiving seed produced in field treated with this product. A copy of this labeling is required to be provided to the conditioner by the producer. All seed alfalfa treated with BIFENTURE EC must be labeled "NOT FOR HUMAN OR ANIMAL CONSUMPTION" at the processing plant. The processor must dispose of all seed screenings in such a way that they cannot be distributed or used for food or feed.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Restricted Entry Interval

12 hours

Timings

[N.A.](#)