

TREE CROPS

General Information

GENERAL INFORMATION

CaB is designed as a foliar nutritional spray to supply Calcium and Boron in a convenient, soluble form. CaB is available and absorbed on the leaf surface or by the root system.

COMPATIBILITY: CaB is compatible with most pesticides. In mixing, always add this product to the tank last. A compatibility test is recommended if the desired combination has not been used in the past.

DIRECTIONS

For best results, spray when the crop is in an active growing state after irrigation or natural rain fall. Spray early in the morning or late afternoon. Midday sprays may not be effective because of excessive moisture evaporation.

CAUTION:

Excessive amounts of Boron may cause damage to susceptible crops. Determine plant sensitivity prior to use. Contact your local agricultural authority for information on specific crops.

Recommendations for Use: CaB may be used on the following crops: Alfalfa, Almonds, Avocados, Apples, Beets, Broccoli, Brussels Sprouts, Cabbage, Carrots, Cauliflower, Celery, Citrus, Clover, Corn, Cotton, Grapes, Lettuce, Milo, Melons, Nectarines, Rice, Pears, Peaches, Pecans, Peppers, Plums, Prunes, Potatoes, Peanuts, Sorghum, Sugar Beets, Sweet Corn, Sugarcane, Strawberries, Tobacco, Tomatoes, Turnips, Walnuts, Watermelons, Wheat and most other crops.

Limitations, Restrictions, and Exceptions

Tree Crops:

Apply 1 to 2 pints per 100 gallons of spray. Start two to three weeks after full bloom and continue at two-week intervals. Trees must be wet thoroughly. For best results do not spray in the heat of the day.

Note: Do not exceed 1 gallon per acre per application.

Foliar Applications:

Conventional Sprayers: Use a minimum of 20 gallons of water per acre.

Low Volume Sprayers: Use 5 gallons of water per acre. If less water is used, slight burning of the foliage may occur.

Aerial Applications: Do not exceed 1 quart per gallon of water.

Method

[Broadcast/Foliar Air](#)

[Spray](#)

[Broadcast/Foliar Air](#)

[Spray](#)

Timings

[Start two to three weeks after full bloom and continue at two week intervals.](#)

[When the crop is in an active growing state after irrigation or natural rain fall.](#)