

## **WHEAT (FORAGE, GRAIN, HAY, STRAW) - OREGON, WASHINGTON AND IDAHO**

### General Information

#### RESTRICTIONS

- This product is solely for use on agricultural crops grown for commercial or research purposes.
- This product may not be applied to residential sites (i.e., in yards, gardens, around homes and apartments) and other sites such as the exterior areas around schools and daycare facilities, and athletic fields, playgrounds, parks, recreation areas, etc.
- Broadcast applications and applications of this product directed to plant parts are prohibited except to grass (grown for seed and/or forage and hay), corn and soybeans, wheat, and clover (grown for seed).
- When broadcast applications are allowed, apply using standard broadcast spreaders.
- Do not allow bait pellet to come in contact with plant parts.
- Use only as a baiting treatment to be applied directly to the soil area beneath listed plants.
- Use as band treatment between rows.
- Grower is responsible for residues on crops.
- Do not apply to dry soil. Make applications following irrigation.
- Do not re-water for 48 hours after application.
- Not for use in California on clover grown for seed.
- Do not use in or around Hanalei, Kaua'i (USFW Refuge)

OR-CAL DUO is a soil surface treatment for control of snails and slugs in citrus groves, berry crops, listed vegetable crops, ornamentals for production, and seed

grasses. When applied as directed, OR-CAL DUO will withstand rain and sprinkling and will offer protections to gardens for several weeks. This product may be injurious to Daylilies and Clematis. Do not use on these plants.

For use on ornamentals and listed crops (other than citrus, grass grown for seed and/or forage and hay, wheat, corn and soybeans), apply in a band, between rows and/or directed towards base of the plants, by hand or with commercial application equipment designed for banded application of granular products.

For use on citrus, apply in a band under tree canopy, by hand or with commercial application equipment designed for banded application of granular products.

For grass (grown for seed and/or forage and hay) and wheat apply by broadcast application.

For clover (grown for seed): Apply as a perimeter treatment at planting and for subsequent applications by broadcast, directing the product to the soil. Not for use in California.

For corn and soybeans: Apply as a perimeter treatment at planting and for subsequent applications by broadcast, directing the product to the soil.

For clover (forage and hay), apply as a perimeter treatment.

Apply any time of day, but evening applications are preferred.

#### Limitations, Restrictions, and Exceptions

Minimum Days Between Retreatment: 14

Preharvest Interval: Prior to formation of seed head

#### Method

[Broadcast](#)

[Soil Surface Treatment](#)

[Bait](#)

Restricted Entry Interval

24 hours

#### Timings

[N.A.](#)