

FOR THE CONTROL OF SLUGS AND SNAILS IN CHRISTMAS TREE PLANTATIONS AND CHRISTMAS TREE LANDING SITES

General Information

RESTRICTIONS

- This product is solely for use on agricultural crops grown for commercial or research purposes, and in State and/or Federal invasive mollusk eradication operations.
- This product may not be applied to residential sites (i.e., in yards, gardens, around homes and apartments) and other sites such as the exterior areas around schools and daycare facilities, and athletic fields, playgrounds, parks, recreation areas, etc.
- Broadcast applications and applications of this product directed to plant parts are prohibited except to grass (grown for seed and/or forage and hay), taro, and as specified for use in State and/or Federal invasive mollusk eradication operations.
- Clover (grown for seed): Apply as a perimeter treatment at planting and for subsequent applications by broadcast, directing the product to the soil.
- Do not allow bait pellet to come in contact with plant parts.
- Use only as a baiting treatment to be applied directly to the soil area beneath listed plants.
- Use as band treatment between rows.
- Do not apply as a wide-area broadcast.
- Grower is responsible for residues on crops.
- Do not apply to dry soil. Make applications following irrigation.
- Do not re-water for 48 hours after application.
- Do not use in or around Hanalei, Kaua'i (USFW Refuge).

Slugger Ultra 4.0 Slug & Snail Bait is a soil surface treatment for control of snails

and slugs in citrus groves, berry crops, listed vegetable crops, ornamental plants, and grass (grown for seed and/or forage and hay).

When applied as directed, Slugger Ultra 4.0 Slug & Snail Bait will withstand rain and sprinkling and will offer protection to gardens for several weeks. This product may be injurious to Daylilies and Clematis. Do not use on these plants. Apply any time of day, but evening applications are preferred.

For use on ornamental plants, artichokes, listed berries, strawberries, celery, rhubarb, Swiss chard, Chinese celery, celtuce, cardoon, Florence fennel, ginseng, beans, peas, prickly pear cactus, tomatoes, lettuce, Cole crops and listed leafy greens and mint apply in a band, between rows and/or directed towards base of the plants, by hand or with commercial application equipment designed for banded application of granular products.

For use on citrus, apply in a band under tree canopy, by hand or with commercial application equipment designed for banded application of granular products.

For grass (grown for seed and/or forage and hay) and taro, apply by broadcast application.

For clover (grown for seed), apply as a perimeter treatment at planting and then for the second application, by broadcast, directing the product to the soil.

For clover (forage and hay), apply as a perimeter treatment.

Limitations, Restrictions, and Exceptions

For the Control of Slugs and Snails in Christmas Tree Plantations and Christmas Tree Landing Sites

INFORMATION

The primary purpose of this SLN is to eliminate slug/snail populations on exported and/or commercially grown harvested Christmas trees. This bait label allows pre-harvest applications to Christmas Tree Plantations. It also allows treatment to landing sites and to fencerows, roadsides, and general non-crop farmstead areas within 100 feet of landing sites.

For best results, in infested areas, begin treatments one year prior to harvesting or cutting. Product should not be applied to dry soil. Applications should be made

following a rain for best results. Bait may be applied during any time of the day, but evening applications are generally better.

DIRECTIONS FOR USE

Christmas Tree (Spring Treatment): Apply before spring rains subside, approximately April 15th to May 30th. When using ground equipment, broadcast bait to the soil area between trees and as far under tree skirts as possible. Apply 15 lbs. to 25 lbs. per acre every 14 to 21 days. Repeat treatment up to 3 applications if slug activity is evident. For spring application this product may be applied aerially, and can be combined with fertilizer.

Christmas Tree (Fall Treatment): Starting on October 1st or after first fall rains, broadcast bait to the soil area between trees and as far under tree skirts as possible. Apply 15 lbs. to 25 lbs. per acre every 14 to 21 days. The last treatment should be made at least 7 days before harvest or cutting. Up to 3 treatments may be necessary depending on slug severity. Apply by ground application only. Do not apply aerially in the fall. Do not apply bait in such a manner that the bait becomes physically lodged in the tree canopy or needles.

Slugger ULTRA 4.0 Slug & Snail Bait may also be used to control slugs and snails at landing sites or in areas in which Christmas trees are baled, stored, handled and/or grouped for shipping. Vegetated fencerows, roadsides, and general non-crop farmstead areas can serve as a source of infestation; these areas within 100 feet of landing sites can be treated. Apply 15 lbs. to 25 lbs. per acre every 14 to 21 days. For smaller areas, use 1/3 lb. to 1/2 lb. to treat 1,000 sq. ft. of soil surface.

For the best efficacy the last treatment should be made at least 7 days before harvest. The entire slug population is not above ground every night to feed. A full feeding cycle is about one week.

RESTRICTION

Do not place bait inside shipping containers.

Do not apply bait directly to trees once they have been cut.

Method

[Broadcast/Foliar Ground](#)

[Bait](#)

[Broadcast/Foliar Ground](#)

Bait

Restricted Entry Interval

12 hours

Timings

(Fall Treatment): Starting on October 1st or after first fall rains.

(Fall Treatment): Starting on October 1st or after first fall rains.