

# **DRY POULTRY LITTER APPLICATIONS**

## General Information

AGROTAIN PLUS SC for UAN nitrogen stabilizer is a liquid suspension additive for urea ammonium nitrate (UAN) solution, liquid manure and dry poultry litter. It contains both a urease inhibitor and nitrification inhibitor suspended in a urea solution. Together these components reduce nitrogen loss from leaching, denitrification and volatilization to allow more complete nitrogen utilization by the crop.

## GENERAL PRECAUTIONS AND DIRECTIONS FOR USE

**WARNING:** Keep out of the reach of children. Pregnant or breastfeeding women must not handle this product. Read the entire label and safety data sheet before use. Use only according to label directions. Always wear proper personal protection when handling the product (long sleeves and long pants, goggles and chemical resistant gloves). Use only with adequate ventilation (large volumes of air movement). If organic vapors or dusts are present above their regulatory limits, follow a respiratory protection program that meets applicable federal, state, provincial or local requirements whenever workplace conditions warrant respirator use.

Do not apply directly to water. Do not contaminate waterways and sewers during cleaning of equipment or disposal of wastes. Always adhere to applicable provisions of federal, state, provincial and local laws and regulations.

Do not use in livestock feed. Keep livestock off treated areas until after 14 days or 1 inch of rainfall has accumulated. UAN may be toxic to livestock when ingested. Do not enter treated areas until spray has thoroughly dried.

## COMPATIBILITY

Once this product is added into the UAN solution, test compatibility with a jar test or similar method before adding other products to the mixture. After a successful test, add other pesticides or surfactants according to the additional product's label.

## Limitations, Restrictions, and Exceptions

## DRY POULTRY LITTER APPLICATIONS:

Don proper personal protection. Add the appropriate amount of product as identified in these directions and mix thoroughly with the litter prior to loading. To ensure adequate coverage of the product on the litter, apply using a sprayer prior to cleaning the hose or loading. Apply only when conducting a complete litter cleanout. Do not apply when animals are present. Do not allow animals to re-enter until after all of the litter has been removed. Apply treated litter to the field immediately. Keep livestock off treated areas until after 14 days or 1 inch of rainfall has accumulated for both liquid and dry manure applications.

Method

[Spray](#)

Timings

[N.A.](#)