

FOR ACTIVE AND HIBERNATING STAGES OF BOXELDER BUGS, ELM LEAF BEETLES, CLUSTER FLIES AND CLOVER MITES - FOR CLOSED VOIDS

General Information

HOW TO USE BASF CRACK & CREVICE INJECTION SYSTEM

CRACK & CREVICE TREATMENTS: Use only with the supplied actuator and injection tubes. Inject into cracks and crevices or void spaces where insects may be harboring, living and breeding. Place injector tip into cracks, crevices, holes and other small openings. Release approximately 1 sec of product. For light infestations, move injector tip along cracks while treating at the rate of 3 ft/sec. For heavy infestations, move injector tip along at 1 ft/sec. For closed voids, calculate the void's cubic area and treat at the rate of 5 to 10 sec/3 ft³. Several holes may be required in long running voids.

SPOT TREATMENTS: To make a spot treatment, release product onto the surface. DO NOT exceed 2 ft² for each spot treatment.

Limitations, Restrictions, and Exceptions

INDOOR AND OUTDOOR TREATMENTS

FOR ACTIVE AND HIBERNATING STAGES OF BOXELDER BUGS, ELM LEAF BEETLES, CLUSTER FLIES AND CLOVER MITES: Inject cracks and crevices and/or voids where these insects may be harboring or hibernating. Infested areas include, but are not limited to, cracks and crevices, behind siding, wall voids, around window and door frames.

Method

[Spot treatment](#)

[Injection](#)

Timings

[N.A.](#)