

FOOD HANDLING ESTABLISHMENTS - FOR LIGHT INFESTATION

General Information

HOW TO USE BASF CRACK & CREVICE INJECTION SYSTEM

CRACK & CREVICE TREATMENTS: Use only with the supplied actuator and injection tubes. Inject into cracks and crevices or void spaces where insects may be harboring, living and breeding. Place injector tip into cracks, crevices, holes and other small openings. Release approximately 1 sec of product. For light infestations, move injector tip along cracks while treating at the rate of 3 ft/sec. For heavy infestations, move injector tip along at 1 ft/sec. For closed voids, calculate the void's cubic area and treat at the rate of 5 to 10 sec/3 ft³. Several holes may be required in long running voids.

SPOT TREATMENTS: To make a spot treatment, release product onto the surface. **DO NOT** exceed 2 ft² for each spot treatment.

Limitations, Restrictions, and Exceptions

FOOD HANDLING ESTABLISHMENTS

Applications are permitted in both food/feed and non-food/feed areas of food/feed handling establishments as spot or Crack & Crevice treatments. Food/feed handling establishments are defined as places other than private residences in which exposed food/feed is held, processed, prepared or served. Food/feed areas are areas for receiving, storing, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups) of food. Serving areas where food is exposed and the facility is in operation are also considered food areas.

SPOT AND CRACK & CREVICE APPLICATION: Spot or Crack & Crevice applications may be made while the facility is in operation, except in federally inspected meat and poultry plants. Cover or remove food from the area being treated. **DO NOT** apply directly to food or food handling surfaces. May be used in federally inspected meat and poultry plants.

Method

[Spot treatment](#)

[Injection](#)

Timings

N.A.