

# **OUTDOOR TREATMENTS - SUBTERRANEAN ANTS SUCH AS ARGENTINE ANTS & HARVESTER ANTS**

## General Information

### HOW TO USE BASF CRACK & CREVICE INJECTION SYSTEM

**CRACK & CREVICE TREATMENT:** Use with the supplied actuator and injection tubes or other BASF equipment. Inject into cracks and crevices or void spaces where insects may be harboring, living and breeding. Place injector tip into cracks, crevices, holes and other small openings. Release approximately 1 sec of product. For light infestations, move injector tip along cracks while treating at the rate of 3 In ft/sec. For heavy infestations, move injector tip along at 1 In ft/sec. For closed voids, calculate the void's cubic area and treat at the rate of 1 to 5 sec/3ft<sup>3</sup>. Several holes may be required in long-running voids.

**SPOT TREATMENT:** Use with the supplied actuator and injection tubes or other BASF equipment. Hold container upright while spraying. Direct nozzle approximately 12" from surface to be treated. Apply directly on insects in these locations when possible.

## Limitations, Restrictions, and Exceptions

### OUTDOOR TREATMENTS

**SUBTERRANEAN ANTS SUCH AS ARGENTINE ANTS and HARVESTER ANTS:** Inject the nest at 3 to 6 sites at a rate of 5 to 10 sec/site. For very large nests (over 36" diameter), increase the number of injection sites. Space injection sites in a circular pattern on the nest surface with 1 site in the center. To quiet active ants on nest surface, sweep surface with spray from a distance of 12". For best results, apply when temperature is 65 to 85°F, or in early morning or late evening hours. Treat new nests as they appear.

## Method

[Injection](#)

Timings

[N.A.](#)