

# **CANEBERRIES (BLACKBERRY, LOGANBERRY, RED AND BLACK RASPBERRY)**

## General Information

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a manner that will contact workers or other persons, either directly or through drift. Only protected workers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency representative responsible for pesticide regulation.

### HOW TO APPLY

Spread the bait on soil prior to planting, prior to plant emergence, and after plant emergence around or near the plants to be protected. For broadcast application, use standard broadcast spreaders. For row application, use standard granular spreaders. Apply the higher rates if the infestation is severe or if the area is heavily watered or after long periods of heavy rain. Reapply as the bait is consumed or at least every 2-4 weeks. Do not apply more than 3 times in any 30 day period.

Do not place in piles. The product works best when soil is moist but with little or no standing water. See specific directions for different use sites.

BRANDT ANTIXX PLUS contains 0.0007 lbs of spinosad per lb. Use the following table to assist in calculating applied lbs spinosad per acre.

See table in the label for APPLICATION RATE REFERENCE.

## Limitations, Restrictions, and Exceptions

### BERRIES:

Spread the bait around the perimeter of the field or greenhouse to intercept pests migrating toward the berries. Use a rate of 20-44 lbs. per acre (0.5-1 lb per 1,000 square feet) and scatter by hand or with granular spreaders. If pests are already in the plots, then carefully spread bait between the furrows near the base of the plants.

### Maximum number of applications

- Do not make more than 6 applications per calendar year.

### Application Interval

- Do not make applications less than 5 days apart.

### Additional Restrictions

- Do not apply more than a total of 0.45 lb ai spinosad (including foliar uses) per acre per crop.

### Method

[Bait](#)

### Pre-Harvest Interval

1 day

### Restricted Entry Interval

4 hours

### Timings

[N.A.](#)