

CONTROL OF CUTWORMS AND LYGUS INFESTING CARROTS GROWN FOR SEED - OREGON

Limitations, Restrictions, and Exceptions

Application Directions

Apply 1 quart of Lorsban Advanced per acre in at least 10 gallons of water per acre as a foliar broadcast spray to ensure thorough coverage of treated areas. For aerial application, apply in a minimum of 5 gallons of water per acre. Apply when field counts or crop injury indicate that damaging pest populations are developing or present. Consult your local cooperative agricultural extension service for additional information concerning cutworm and lygus control practices in your area.

Specific Use Restrictions:

- Maximum single application rate is 0.94 lb ai chlorpyrifos (2 pints of Lorsban Advanced) per acre.
- Do not make more than one application of Lorsban Advanced or any other product containing chlorpyrifos per crop per year.
- This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Evening or night time applications when foragers are not present reduce exposure hazards. Consult your cooperative agricultural extension service for additional information concerning protection of pollinating bees.
- Lorsban Advanced should not be tank mixed with other pesticides, surfactants, or fertilizer formulations unless prior use experience has shown that combination to be noninjurious under local conditions of use.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Timings

[When field counts or crop injury indicate that damaging pest populations are developing or present.](#)