

# **TUBEROUS AND CORM VEGETABLES (SUBGROUP 1C) - WEEDS CONTROL AT 1.0-2.0 FL OZ PER ACRE**

## General Information

### PRODUCT INFORMATION

AIM EC is an emulsifiable concentrate formulation. AIM EC is to be mixed with water, liquid fertilizer or mixtures of water and liquid fertilizer and adjuvants and applied to labeled crops and non-crop areas for selective postemergence control of broadleaf weeds, for sucker control, for burndown prior to planting, as a harvest aid and to defoliate/desiccate labeled crops.

Weed control is optimized when the product is applied to actively growing weeds. AIM EC is a contact herbicide. Within a few hours following application, the foliage of susceptible weeds show signs of desiccation. Extremes in environmental conditions such as temperature, moisture, soil conditions, and cultural practices may affect the activity of AIM EC. Herbicide symptoms may be accelerated under moist conditions. Weed control may be reduced when weeds are hardened off by drought and become less susceptible to AIM EC.

AIM EC is rapidly absorbed through the foliage of plants. To avoid significant crop response, applications should not be made within 6 to 8 hours of either rain or irrigation or when heavy dew is present on the crop. Environmental conditions and certain spray tank additives may increase herbicidal symptoms on the crop.

### ADJUVANT USE REQUIREMENTS

The use of a quality spray adjuvant is required for optimum performance. Refer to the individual crop recommendation sections of this label for specific adjuvant type and use rates.

### ON-FARM TESTING

Not all varieties or cultivars of labeled crops have been fully evaluated under all environmental and soil conditions. Consult with your local seed company for additional information.

It may also be beneficial to conduct small on-farm trials under actual conditions with specific varieties or cultivars before treating large acreage.

## APPLICATION INFORMATION

### APPLICATION METHODS

#### GROUND APPLICATION

Use ground sprayers designed, calibrated and operated to deliver uniform spray droplets to the targeted plant or plant parts. Adjust sprayer nozzles to achieve uniform plant coverage. Overlaps and slower ground speeds (caused by continuing to spray while starting, stopping or turning) may result in higher application rates and possible crop response.

#### Spray Buffer for Ground Application

Spray buffer zones for ground applications, listed in table below, are required near desirable perennial vegetation or crops before blossom and after total leaf drop, and/or near other desirable or annual crops.

#### Broadcast Boom Sprayers

Use a broadcast boom sprayer equipped with the appropriate nozzles, spray tips and screens and adjusted to provide optimum spray distribution and coverage at the appropriate operating pressures. Use nozzles that produce minimal amounts of fine spray droplets. Do not exceed 30 psi spray pressure unless otherwise required by the manufacturer of drift reducing nozzles. Apply a minimum of 10 gallons of finished spray per acre. Use higher spray volumes when there is a dense weed population or crop canopy. Adjust sprayers to position spray tips no lower than 12-18 inches above the crop or weed canopy depending on the nozzle specification. Operate the sprayer to avoid the application of high herbicide rates directly over the rows or into the whorl of treated crop plants.

#### Directed Sprayers

For directed sprayers apply AIM EC with drop nozzles or other spray equipment.

Post directed applications may be utilized when labeled crops have reached minimum growth stages where sprays may be directed to the target weeds, but is not deposited on the green stem, foliage, blooms or fruit of the crop. Do not apply when conditions favor drift or when is above 10MPH.

Use drop nozzles or other spray equipment capable of directing the spray to target weeds and away from sensitive plant parts. Apply when labeled crops have reached minimum growth stages described in specific crop sections of this label and when spray will not be deposited on green stems, foliage, blossoms or fruit.

#### Hooded Sprayers

To apply AIM EC using a hooded sprayer, refer to the Hooded Sprayer Section for specific adjustment and operation instructions. For additional information, refer to the individual crop sections of this label.

#### Hand held or high volume orchard gun sprayers

AIM EC may be applied to certain labeled crops and non-crop areas with hand operated sprayers such as backpack sprayers, compression sprayers, knapsack sprayers, or high volume orchard gun sprayers. Directed applications may be utilized when labeled crops have reached minimum growth stages where sprays may be directed to the target weeds, but is not deposited on the green stem, foliage, blooms or fruit of the crop. Refer to individual crop sections of this label.

#### AERIAL APPLICATION

Use nozzle types and arrangements that will provide optimum coverage while producing a minimal amount of fine droplets. Apply at a minimum of 3 gallons of finished spray per acre. Spray volumes greater than 3 GPA may be needed for harvest aid and defoliation treatments, or for dense weed populations or with dense crop canopies.

#### CROP ROTATIONAL RESTRICTIONS

Following an application of AIM EC, a treated field may be rotated to a registered crop at any time, subject to specific crop restrictions that may be found in the individual crop sections. All other crops may be planted after 12 months.

#### Limitations, Restrictions, and Exceptions

#### TUBEROUS AND CORM VEGETABLES (SUBGROUP 1C)

##### Restrictions

Do not apply more than 11.6 fl oz of (0.181 pound active ingredient) per acre per crop season as a desiccant.

Do not apply when conditions favor drift or wind is above 10 mph.

## DIRECTIONS FOR USE

Apply AIM EC alone or in a tank mix combination with other herbicides and insecticides as a fallow systems treatment, as a preplant burndown treatment and/or as a harvest aid to desiccate potatoes and those susceptible weeds that may be present.

### Preplant Burndown

Apply AIM EC alone or with other herbicides or liquid fertilizers as a burn-down treatment to control or suppress weeds. AIM EC is effective as a burndown treatment for crops prior to new plantings. Do not exceed the applicable amounts as listed for the specific crop in the MAXIMUM ALLOWABLE AIM EC USE in Table 1. For optimum performance, make applications to actively growing weeds up to 4 inches high or rosettes less than 3 inches across. Coverage is essential for good control. Optimum broad-spectrum control of annual and perennial weeds requires a tank mix with a labeled burndown herbicides such as glyphosate, glufosinate, paraquat, 2,4-D, or dicamba.

### Ground Application

Apply AIM EC in at least 20 gallons of water per acre. Vary the spray volume and spray pressure as indicated by the density of the potato canopy and vines to assure thorough spray coverage. Increase the spray volume and pressure if the potato canopy is dense or under cool, cloudy or dry conditions. Increased spray volumes will enhance performance.

### Aerial Application

Apply AIM EC with aerial equipment using 5 to 10 gallons of water per acre, using higher volumes when potato canopies and vines are dense. Adjust the nozzles to provide a uniform pattern and a droplet size of 350 to 450 microns.

### Adjuvant Requirements

A nonionic surfactant (NIS), methylated seed oil (MSO), or crop oil concentrate (COC) or other suitable surfactant mixture is required. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient, or a methylated seed oil, or crop oil concentrate (COC)(petroleum or seed oil) at 1 to 2 v/v (1 to 2 gallons per 100 gallons of spray solution. The use of a high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per

100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the nonionic surfactant methylated seed oil or crop oil is allowed.

Adjuvant rates should increase as spray volumes exceed 20 gallons per acre.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

7 days

Restricted Entry Interval

12 hours

Timings

[Preplant](#)