

# **DRIED SHELLLED BEANS, PEAS CROP SUBGROUP 6-C: AND FLAX (EXCEPT SOYBEAN) AND VEGETABLE FOLIAGE OF LEGUME GROUP 7 - HARVEST AID APPLICATION**

## General Information

### PRODUCT INFORMATION

AIM EC is an emulsifiable concentrate formulation. AIM EC is to be mixed with water, liquid fertilizer or mixtures of water and liquid fertilizer and adjuvants and applied to labeled crops and non-crop areas for selective postemergence control of broadleaf weeds, for sucker control, for burndown prior to planting, as a harvest aid and to defoliate/desiccate labeled crops.

Weed control is optimized when the product is applied to actively growing weeds. AIM EC is a contact herbicide. Within a few hours following application, the foliage of susceptible weeds show signs of desiccation. Extremes in environmental conditions such as temperature, moisture, soil conditions, and cultural practices may affect the activity of AIM EC. Herbicide symptoms may be accelerated under moist conditions. Weed control may be reduced when weeds are hardened off by drought and become less susceptible to AIM EC.

AIM EC is rapidly absorbed through the foliage of plants. To avoid significant crop response, applications should not be made within 6 to 8 hours of either rain or irrigation or when heavy dew is present on the crop. Environmental conditions and certain spray tank additives may increase herbicidal symptoms on the crop.

### ADJUVANT USE REQUIREMENTS

The use of a quality spray adjuvant is required for optimum performance. Refer to the individual crop recommendation sections of this label for specific adjuvant type and use rates.

### ON-FARM TESTING

Not all varieties or cultivars of labeled crops have been fully evaluated under all environmental and soil conditions. Consult with your local seed company for additional information.

It may also be beneficial to conduct small on-farm trials under actual conditions

with specific varieties or cultivars before treating large acreage.

## APPLICATION INFORMATION

### APPLICATION METHODS

#### GROUND APPLICATION

Use ground sprayers designed, calibrated and operated to deliver uniform spray droplets to the targeted plant or plant parts. Adjust sprayer nozzles to achieve uniform plant coverage. Overlaps and slower ground speeds (caused by continuing to spray while starting, stopping or turning) may result in higher application rates and possible crop response.

#### Spray Buffer for Ground Application

Spray buffer zones for ground applications, listed in table below, are required near desirable perennial vegetation or crops before blossom and after total leaf drop, and/or near other desirable or annual crops.

#### Broadcast Boom Sprayers

Use a broadcast boom sprayer equipped with the appropriate nozzles, spray tips and screens and adjusted to provide optimum spray distribution and coverage at the appropriate operating pressures. Use nozzles that produce minimal amounts of fine spray droplets. Do not exceed 30 psi spray pressure unless otherwise required by the manufacturer of drift reducing nozzles. Apply a minimum of 10 gallons of finished spray per acre. Use higher spray volumes when there is a dense weed population or crop canopy. Adjust sprayers to position spray tips no lower than 12-18 inches above the crop or weed canopy depending on the nozzle specification. Operate the sprayer to avoid the application of high herbicide rates directly over the rows or into the whorl of treated crop plants.

#### Directed Sprayers

For directed sprayers apply AIM EC with drop nozzles or other spray equipment.

Post directed applications may be utilized when labeled crops have reached minimum growth stages where sprays may be directed to the target weeds, but is not deposited on the green stem, foliage, blooms or fruit of the crop. Do not apply when conditions favor drift or when is above 10MPH.

Use drop nozzles or other spray equipment capable of directing the spray to target weeds and away from sensitive plant parts. Apply when labeled crops have reached

minimum growth stages described in specific crop sections of this label and when spray will not be deposited on green stems, foliage, blossoms or fruit.

#### Hooded Sprayers

To apply AIM EC using a hooded sprayer, refer to the Hooded Sprayer Section for specific adjustment and operation instructions. For additional information, refer to the individual crop sections of this label.

#### Hand held or high volume orchard gun sprayers

AIM EC may be applied to certain labeled crops and non-crop areas with hand operated sprayers such as backpack sprayers, compression sprayers, knapsack sprayers, or high volume orchard gun sprayers. Directed applications may be utilized when labeled crops have reached minimum growth stages where sprays may be directed to the target weeds, but is not deposited on the green stem, foliage, blooms or fruit of the crop. Refer to individual crop sections of this label.

#### AERIAL APPLICATION

Use nozzle types and arrangements that will provide optimum coverage while producing a minimal amount of fine droplets. Apply at a minimum of 3 gallons of finished spray per acre. Spray volumes greater than 3 GPA may be needed for harvest aid and defoliation treatments, or for dense weed populations or with dense crop canopies.

#### CROP ROTATIONAL RESTRICTIONS

Following an application of AIM EC, a treated field may be rotated to a registered crop at any time, subject to specific crop restrictions that may be found in the individual crop sections. All other crops may be planted after 12 months.

#### Limitations, Restrictions, and Exceptions

Dried shelled beans, peas Crop Subgroup 6-C: and flax (except soybean) and vegetable foliage of legume Group 7

#### DIRECTIONS FOR USE

##### Harvest AID Treatment:

Apply Aim EC Herbicide as a harvest aid to dry beans and dry peas at maturity when 80 to 90% of seed pods are yellow or buck skin in color and only 30% of green leaves remain on the plant. Apply to flax when 75% of the bolls have turned brown. Thorough coverage is essential for harvest aid and multiple applications may be

needed. For optimum performance use 15 to 30 gallons per acre finished sprayed with a methylated seed oil (MSO) type adjuvant to ensure thorough coverage and retention for harvest aid.

#### AIM EC Use rates:

Apply Aim EC alone or as a tank mixture with other harvest aids. Apply Aim EC at 1.0 to 6.1 fl oz (0.016 to 0.096 pounds active ingredient) per acre, but not to exceed maximum labeled rates. Refer to the MAXIMUM ALLOWABLE AIM USE RATE CHART and the PREHARVEST INTERVAL charts for additional application information.

Applications shall be made in spray volumes sufficient to provide complete coverage of foliage. Use a minimum of 15 gallons of finished spray per acre for ground application and 5 gallons per acre for aerial application. A methylated seed oil (MSO) or crop oil concentrate (COC) is required at 1 to 2 v/v (1 to 2 gallons per 100 gallons of spray solution). The addition of a high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the methylated seed oil or crop oil may enhance performance.

If spraying dry beans before full maturity and pods are not all mature and turning color, a repeat application may be necessary.

#### Note

If applied as a tank mixture, refer to the other product's label for restrictions on tank mixing, and observe all label precautions, instructions and rotational cropping restrictions.

#### Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Restricted Entry Interval

12 hours

#### Timings

[To dry beans and dry peas: At maturity when 80 to 90% of seed pods are yellow or buck skin in color and only 30% of green leaves remain on the plant.](#)

To dry beans and dry peas: At maturity when 80 to 90% of seed pods are yellow or buck skin in color and only 30% of green leaves remain on the plant.