

TOBACCO

General Information

PRODUCT INFORMATION

AIM EC is an emulsifiable concentrate formulation. AIM EC is to be mixed with water, liquid fertilizer or mixtures of water and liquid fertilizer and adjuvants and applied to labeled crops and non-crop areas for selective postemergence control of broadleaf weeds, for sucker control, for burndown prior to planting, as a harvest aid and to defoliate/desiccate labeled crops.

Weed control is optimized when the product is applied to actively growing weeds. AIM EC is a contact herbicide. Within a few hours following application, the foliage of susceptible weeds show signs of desiccation. Extremes in environmental conditions such as temperature, moisture, soil conditions, and cultural practices may affect the activity of AIM EC. Herbicide symptoms may be accelerated under moist conditions. Weed control may be reduced when weeds are hardened off by drought and become less susceptible to AIM EC.

AIM EC is rapidly absorbed through the foliage of plants. To avoid significant crop response, applications should not be made within 6 to 8 hours of either rain or irrigation or when heavy dew is present on the crop. Environmental conditions and certain spray tank additives may increase herbicidal symptoms on the crop.

ADJUVANT USE REQUIREMENTS

The use of a quality spray adjuvant is required for optimum performance. Refer to the individual crop recommendation sections of this label for specific adjuvant type and use rates.

ON-FARM TESTING

Not all varieties or cultivars of labeled crops have been fully evaluated under all environmental and soil conditions. Consult with your local seed company for additional information.

It may also be beneficial to conduct small on-farm trials under actual conditions with specific varieties or cultivars before treating large acreage.

APPLICATION INFORMATION

APPLICATION METHODS

GROUND APPLICATION

Use ground sprayers designed, calibrated and operated to deliver uniform spray droplets to the targeted plant or plant parts. Adjust sprayer nozzles to achieve uniform plant coverage. Overlaps and slower ground speeds (caused by continuing to spray while starting, stopping or turning) may result in higher application rates and possible crop response.

Spray Buffer for Ground Application

Spray buffer zones for ground applications, listed in table below, are required near desirable perennial vegetation or crops before blossom and after total leaf drop, and/or near other desirable or annual crops.

Broadcast Boom Sprayers

Use a broadcast boom sprayer equipped with the appropriate nozzles, spray tips and screens and adjusted to provide optimum spray distribution and coverage at the appropriate operating pressures. Use nozzles that produce minimal amounts of fine spray droplets. Do not exceed 30 psi spray pressure unless otherwise required by the manufacturer of drift reducing nozzles. Apply a minimum of 10 gallons of finished spray per acre. Use higher spray volumes when there is a dense weed population or crop canopy. Adjust sprayers to position spray tips no lower than 12-18 inches above the crop or weed canopy depending on the nozzle specification. Operate the sprayer to avoid the application of high herbicide rates directly over the rows or into the whorl of treated crop plants.

Directed Sprayers

For directed sprayers apply AIM EC with drop nozzles or other spray equipment.

Post directed applications may be utilized when labeled crops have reached minimum growth stages where sprays may be directed to the target weeds, but is not deposited on the green stem, foliage, blooms or fruit of the crop. Do not apply when conditions favor drift or when is above 10MPH.

Use drop nozzles or other spray equipment capable of directing the spray to target weeds and away from sensitive plant parts. Apply when labeled crops have reached minimum growth stages described in specific crop sections of this label and when

spray will not be deposited on green stems, foliage, blossoms or fruit.

Hooded Sprayers

To apply AIM EC using a hooded sprayer, refer to the Hooded Sprayer Section for specific adjustment and operation instructions. For additional information, refer to the individual crop sections of this label.

Hand held or high volume orchard gun sprayers

AIM EC may be applied to certain labeled crops and non-crop areas with hand operated sprayers such as backpack sprayers, compression sprayers, knapsack sprayers, or high volume orchard gun sprayers. Directed applications may be utilized when labeled crops have reached minimum growth stages where sprays may be directed to the target weeds, but is not deposited on the green stem, foliage, blooms or fruit of the crop. Refer to individual crop sections of this label.

AERIAL APPLICATION

Use nozzle types and arrangements that will provide optimum coverage while producing a minimal amount of fine droplets. Apply at a minimum of 3 gallons of finished spray per acre. Spray volumes greater than 3 GPA may be needed for harvest aid and defoliation treatments, or for dense weed populations or with dense crop canopies.

CROP ROTATIONAL RESTRICTIONS

Following an application of AIM EC, a treated field may be rotated to a registered crop at any time, subject to specific crop restrictions that may be found in the individual crop sections. All other crops may be planted after 12 months.

Limitations, Restrictions, and Exceptions

TOBACCO

DIRECTIONS FOR USE

Apply AIM EC alone or as a tank mixture with other registered herbicides to emerged and actively growing weeds. For optimum performance, make applications to weeds up to 4 inches tall and rosettes less than 3 inches across. Use higher rates when treating more mature weeds or dense vegetative growth.

Coverage is essential for good control.

Adjuvant Requirements

Use adequate spray volume to achieve thorough coverage, but a minimum of 10 gallons of finished spray per acre is required. Use a quality crop oil concentrate (COC) at 1% v/v (1 gallon of COC per 100 gallons of spray solution).

AIM EC may be tank mixed with other herbicides registered for use on tobacco to provide additional weed control. For specific mixing instructions, refer to the Mixing and Loading Instructions under the PRODUCT INFORMATION section. Refer to the other product label for restrictions on tank mixing and observe all label precautions, instructions and rotational cropping restrictions.

For additional information refer to the PRODUCT INFORMATION section of the AIM EC label.

Pre-transplant burndown

AIM EC is a contact herbicide for pre-transplant burndown control of broadleaf weeds in tobacco. Apply AIM EC as a broadcast application alone or as a tank mixture with other herbicides to emerged and actively growing weeds. Apply AIM EC up to one (1) day prior to transplanting.

Shielded spray or Hooded spray

Apply AIM EC using shielded sprayers or hooded sprayers to emerged and actively growing broadleaf weeds in tobacco from transplanting until layby. Shielded spray or hooded spray applications of AIM EC or AIM EC tank mixtures should utilize application equipment that must prevent contact of spray solution with the tobacco plant. Do not allow spray solution to contact tobacco foliage or green stem tissue. Refer to the Hooded Sprayer Applications section of this label for additional specific use directions.

Directed spray after first priming (Flue Cured Tobacco Only)

Apply AIM EC as a directed spray application after the first priming in only flue cured tobacco only for the control of emerged and actively growing broadleaf weeds. Directed spray equipment should position nozzles a minimum of 3 to 4 inches above the soil, with nozzles directed underneath the crop canopy. Spray solution should be directed at the base of tobacco plants for minimal contact with foliage while maintaining maximum contact with broadleaf weeds that are at appropriate treatment size. Do not apply when conditions favor drift or wind is above 10 mph.

For control of additional broadleaf weeds and grasses, AIM EC may be tank mixed with other herbicides registered for use in tobacco at the appropriate timing. Refer

to weed control list in Table 3 for appropriate weed control information. Refer to the other product's label for restrictions on tank mixing, and observe all label precautions, instructions and rotational cropping restrictions.

Method

[Directed Spray](#)

[Hooded spray](#)

[Shielded spray](#)

[Directed Spray](#)

[Hooded spray](#)

[Shielded spray](#)

Pre-Harvest Interval

6 days

Restricted Entry Interval

12 hours

Timings

[Postemergence \(Weed\)](#)

[Pretransplant](#)