

# **LOW GROWING BERRY SUBGROUP 13-07G - WEEDS CONTROL AT 0.5 FL OZ PER ACRE**

## General Information

### PRODUCT INFORMATION

AIM EC is an emulsifiable concentrate formulation. AIM EC is to be mixed with water, liquid fertilizer or mixtures of water and liquid fertilizer and adjuvants and applied to labeled crops and non-crop areas for selective postemergence control of broadleaf weeds, for sucker control, for burndown prior to planting, as a harvest aid and to defoliate/desiccate labeled crops.

Weed control is optimized when the product is applied to actively growing weeds. AIM EC is a contact herbicide. Within a few hours following application, the foliage of susceptible weeds show signs of desiccation. Extremes in environmental conditions such as temperature, moisture, soil conditions, and cultural practices may affect the activity of AIM EC. Herbicide symptoms may be accelerated under moist conditions. Weed control may be reduced when weeds are hardened off by drought and become less susceptible to AIM EC.

AIM EC is rapidly absorbed through the foliage of plants. To avoid significant crop response, applications should not be made within 6 to 8 hours of either rain or irrigation or when heavy dew is present on the crop. Environmental conditions and certain spray tank additives may increase herbicidal symptoms on the crop.

### ADJUVANT USE REQUIREMENTS

The use of a quality spray adjuvant is required for optimum performance. Refer to the individual crop recommendation sections of this label for specific adjuvant type and use rates.

### ON-FARM TESTING

Not all varieties or cultivars of labeled crops have been fully evaluated under all environmental and soil conditions. Consult with your local seed company for additional information.

It may also be beneficial to conduct small on-farm trials under actual conditions with specific varieties or cultivars before treating large acreage.

## APPLICATION INFORMATION

### APPLICATION METHODS

#### GROUND APPLICATION

Use ground sprayers designed, calibrated and operated to deliver uniform spray droplets to the targeted plant or plant parts. Adjust sprayer nozzles to achieve uniform plant coverage. Overlaps and slower ground speeds (caused by continuing to spray while starting, stopping or turning) may result in higher application rates and possible crop response.

#### Spray Buffer for Ground Application

Spray buffer zones for ground applications, listed in table below, are required near desirable perennial vegetation or crops before blossom and after total leaf drop, and/or near other desirable or annual crops.

#### Broadcast Boom Sprayers

Use a broadcast boom sprayer equipped with the appropriate nozzles, spray tips and screens and adjusted to provide optimum spray distribution and coverage at the appropriate operating pressures. Use nozzles that produce minimal amounts of fine spray droplets. Do not exceed 30 psi spray pressure unless otherwise required by the manufacturer of drift reducing nozzles. Apply a minimum of 10 gallons of finished spray per acre. Use higher spray volumes when there is a dense weed population or crop canopy. Adjust sprayers to position spray tips no lower than 12-18 inches above the crop or weed canopy depending on the nozzle specification. Operate the sprayer to avoid the application of high herbicide rates directly over the rows or into the whorl of treated crop plants.

#### Directed Sprayers

For directed sprayers apply AIM EC with drop nozzles or other spray equipment.

Post directed applications may be utilized when labeled crops have reached minimum growth stages where sprays may be directed to the target weeds, but is not deposited on the green stem, foliage, blooms or fruit of the crop. Do not apply when conditions favor drift or when is above 10MPH.

Use drop nozzles or other spray equipment capable of directing the spray to target weeds and away from sensitive plant parts. Apply when labeled crops have reached minimum growth stages described in specific crop sections of this label and when spray will not be deposited on green stems, foliage, blossoms or fruit.

#### Hooded Sprayers

To apply AIM EC using a hooded sprayer, refer to the Hooded Sprayer Section for specific adjustment and operation instructions. For additional information, refer to the individual crop sections of this label.

#### Hand held or high volume orchard gun sprayers

AIM EC may be applied to certain labeled crops and non-crop areas with hand operated sprayers such as backpack sprayers, compression sprayers, knapsack sprayers, or high volume orchard gun sprayers. Directed applications may be utilized when labeled crops have reached minimum growth stages where sprays may be directed to the target weeds, but is not deposited on the green stem, foliage, blooms or fruit of the crop. Refer to individual crop sections of this label.

#### AERIAL APPLICATION

Use nozzle types and arrangements that will provide optimum coverage while producing a minimal amount of fine droplets. Apply at a minimum of 3 gallons of finished spray per acre. Spray volumes greater than 3 GPA may be needed for harvest aid and defoliation treatments, or for dense weed populations or with dense crop canopies.

#### CROP ROTATIONAL RESTRICTIONS

Following an application of AIM EC, a treated field may be rotated to a registered crop at any time, subject to specific crop restrictions that may be found in the individual crop sections. All other crops may be planted after 12 months.

#### Limitations, Restrictions, and Exceptions

##### LOW GROWING BERRY Subgroup 13-07G

##### Restrictions

Do not apply more than 2 fl oz (0.031 lb ai) during the dormant season.

Do not apply more than 6.15 fl oz per acre per season (0.096 pound active ingredient per acre per season).

## DIRECTIONS FOR USE

AIM EC Herbicide applications will control susceptible emerged broadleaf weeds. Repeat applications may be necessary for weeds that emerge after an AIM EC Herbicide treatment.

Apply only by ground equipment such as boom sprayers, shielded or hooded sprayers, hand-held or high-volume wands or orchard guns. Use a minimum of 20 gallons finished spray solution per broadcast acre.

### Dormant Applications

Apply AIM EC Herbicide as a broadcast application to the base of the trunk to control emerged and actively growing weeds during the dormant stage of the crop.

### Post-directed Applications for Broadleaf Weed Control

Apply AIM EC Herbicide as a directed spray avoiding contact with the berry plant but directed at actively growing weeds. AIM EC Herbicide is a contact herbicide and coverage is essential for good weed control. Do not allow AIM EC Herbicide spray mist to come in contact with green stem tissue, desirable fruit, blooms or foliage. Newly planted bush berries should only be treated with shielded sprayers or hooded sprayers.

### AIM EC Herbicide Use Rates

Apply up to 2 fl oz (0.031 pound active ingredient) AIM EC Herbicide per broadcast acre. For best control, apply to seedling weeds in the 2 to 3-leaf stage. Use higher labeled rates of AIM EC Herbicide for larger weeds up to 6 leaves. Weeds greater than 6 leaves may be only partially controlled. See Table 3 for AIM EC Herbicide use rates and weeds controlled.

## Adjuvant Requirements

A nonionic surfactant (NIS), methylated seed oil (MSO) or crop oil concentrate (COC) is required. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient, or a methylated seed oil, or crop oil concentrate (COC)(petroleum or seed oil) at 1 to 2 v/v (1 to 2 gallons per 100 gallons of spray solution). A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the nonionic surfactant methylated seed oil or crop oil is allowed.

AIM EC Herbicide may be mixed with other registered herbicides for broader spectrum weed control. When tank mixing with fertilizer solutions, be sure to prepare an AIM EC Herbicide premixture of AIM EC Herbicide and clean water. See Mixing and Loading Instructions under the PRODUCT INFORMATION section of this label for specific mixing instructions. Refer to this and the other product's labels for mixing instructions, precautions, and restrictions. Follow the most restrictive instructions for each tank-mix partner.

## Precautions

Extreme caution must be taken during applications when desirable fruit, foliage and/or blooms are present in order to avoid spotting or necrosis. Do not allow AIM EC Herbicide spray mist to come in contact with green stem tissue, desirable fruit, blooms or foliage.

For seedling or newly transplanted bushes, do not allow spray to contact green bark of trunk area. Use shielded sprayers only.

## Method

[Broadcast/Foliar Ground](#)

[Directed Spray](#)

Restricted Entry Interval

12 hours

## Timings

[Postemergence \(Weed\)](#)