

TURF TANK MIXES FOR SUMMER STRESS, ANTHRACNOSE

General Information

PRODUCT INFORMATION

Appear II Fungicide is a systemic fungicide, which contains potassium phosphite. Appear II Fungicide is effective in prevention and control of diseases caused by Pythium, Microdochium, Anthracnose, and Rhizoctonia spp. Appear II Fungicide, applied as part of an Integrated Pest Management (IPM) program with good cultural management, promotes a vigorous, healthy stand of turf that is better able to tolerate stress. In order to achieve maximum disease control, apply before the expression of disease or when disease is first observed. Initiate applications when environmental conditions are favorable for disease development.

MIXING INSTRUCTIONS

Apply in 1-5 gallons of water per thousand square feet. Fill spray tank halfway with clean water. Add the required amount of Appear II Fungicide to the tank as indicated in the tables below. While agitating, add the remaining amount of water.

COMPATIBILITY

Appear II Fungicide is believed to be compatible with most commonly used pesticides. Consult specific product labels for additional information. Conduct a tank spray compatibility test when you plan to mix this product with other products. To determine the physical compatibility of this product with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to approximately one quart of water with agitation. Add dry formulations first, then flowables, and then emulsifiable concentrates last. After thorough mixing, allow this mixture to stand for 5 minutes. If the combination remains mixed or can be readily remixed, it is physically compatible. Once compatibility has been proven, use the same procedure for adding products to the spray tank.

Appear II Fungicide has been evaluated for phytotoxicity on a large number of turf grasses under various growing conditions. However, testing all plant species, with all tank mixtures and combinations is not feasible. Test a small portion of the area to be treated for phytotoxicity, prior to treating the entire area.

Limitations, Restrictions, and Exceptions

Turf Tank Mixes for Control of Summer Stress, Anthracnose, Bermudagrass Decline, Bermudagrass Leaf Spot, and Pink Snow Mold

An Apear II Fungicide program encourages the development a vigorous, healthy stand of turf that is better able to tolerate stress as part of a complete disease control program. Apply Apear II Fungicide with products registered to control the disease complex on turf in accordance with the most restrictive of label limitations and precautions. Do not exceed the label dosage rates.

Tank Mix Product: Daconil Action, Secure

Application Interval: 14 days

Instructions

Make applications prior to the appearance of the disease and reapply on a 14 day interval while conditions favor the disease.

- Use a shorter spray interval and a higher rate under extra disease pressure.

Method

[Spray](#)

Timings

[Prior to the appearance of the disease.](#)