

CEREALS, WHEAT - FOOT ROT/EYESPOT

Limitations, Restrictions, and Exceptions

CEREALS, WHEAT

Remarks

Apply full rate of COVER XL plus half the rate specified on other EPA registered fungicides such as Topsin M. Apply at tillering but before elongation has occurred.

APPLICATION: COVER XL is most effective when applied and allowed to dry before a rainfall. For best results, sufficient coverage is very important. Use a higher water volume for aerial application (greater than 2 GPA) if equipment and/or conditions would not provide good coverage. COVER XL may be applied by ground, air, or chemigation.

WHEAT USE RESTRICTIONS:

1. Do not apply more than 2 applications/A/year.
2. Do not apply after Feekes 10.54.
3. Do not apply more than 28 fl. oz./A/season of COVER XL.
4. Do not apply more than 0.22 lb. a.i. propiconazole-containing products/A/season.
5. Do not apply more than 0.40 lb. a.i. azoxystrobin-containing products/A/season.
6. Under certain environmental conditions, tank mixes of COVER XL plus herbicides and/or fertilizers may cause crop injury.
7. Do not apply within 7 days of harvest (7-day PHI) for forage and hay.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

7 days

Restricted Entry Interval

12 hours

Timings

At tillering but before elongation has occurred.