

GRAIN SORGHUM - AT BEDDING

General Information

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation. This label must be in the possession of the user at the time of pesticide application

PRECAUTIONS AND RESTRICTIONS

BEFORE USING, READ PRECAUTIONARY STATEMENTS AND ENVIRONMENTAL HAZARDS.

- Do not contaminate food or feed products.
- Not for use or storage in or around home.
- Keep out of reach of Domestic Animals.
- Once a container has been attached to the hopper, it is recommended that it be used completely. Make sure the hoppers are emptied while still in the field. Refer to Storage and Disposal statements for further instructions.

Loader and applicator requirements to minimize the potential for runoff to surface water:

- Maintain a 500 ft vegetative buffer between treated area and surface water on neighboring land.
- Maintain a 500 ft buffer between a standpipe drain outlet and surface water on neighboring land.
- Maintain a 66 ft setback between the treated area and entry points to surface water bodies on non-highly erodible soils.
- Maintain a 300 ft setback between the treated area and entry points to surface water bodies on highly erodible soils (HES).
- Maintain a 66 ft setback between treated area and standpipes on terraced fields as well as a 66 ft vegetative buffer between the tile outlet and entry points to surface water bodies.

All setbacks must be planted to a crop or seeded with grass or other suitable cover, and all vegetative buffers must be seeded with grass or other suitable cover.

DO NOT load, rinse, or wash equipment within 300 ft from surface water bodies.

DO NOT load, rinse, or wash equipment within 50 ft from wells unless conducted on an impervious surface.

To prevent channeling of surface water runoff, adjust the planter row-cleaners appropriately to prevent rows lower in height than adjacent soil.

CALIBRATION INFORMATION

It is important that applicator equipment be properly set to deliver the labeled rate. Please refer to planter manufacturer for detailed information on appropriate gauge settings and calibration procedures.

Refer in the label for Use Rates as Pounds per Acre (based on various row spacing and application rates)

Limitations, Restrictions, and Exceptions

GRAIN SORGHUM - At Bedding

Rate

- 5.2 oz. per 1,000 ft. of row for any row spacing (minimum 20 inch row spacing) or no more than 8.4 lbs. of Counter 20G per acre.

Application

- Knifed-in: Drill granules 1-4 inches below the seed OR 1-4 inches below the seed and up to 5 inches to the side of the seed.

Remarks

Restriction: DO NOT place granules in direct contact with seed as crop injury may occur.

Restriction: DO NOT graze or harvest grain sorghum forage within 50 days after treatment.

Restriction: DO NOT harvest grain sorghum fodder or grain within 100 days after treatment.

Method

[Knifed-in](#)

Pre-Harvest Interval

Forage - 50 days

Fodder or Grain - 100 days

Restricted Entry Interval

48 hours

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[At Bedding](#)