

CITRUS AND AVOCADOS, TREE FRUIT AND NUT CROPS

General Information

GENERAL INFORMATION

*Formula 1 S.W.0-28-25 is a foliar nutritional spray which is intended to be used as a supplemental fertilizer treatment. to a regular fertilizer program. Upon foliar application, the phosphite ions are taken up directly by the plant foliage and may undergo a degree of conversion to phosphate ions, or will be used directly by plants as phosphite ions. As a soil application to annual crops, a lesser response from the initial crop, with a corresponding superior response from succeeding crops may be observed. In addition, placement close to seeds or root zones may be injurious to crops. The effect may be aggravated by a soil pH below 6.5. Because of varying conditions over which the manufacturer has no control, including weather, the use of other pesticides and fertilizers and variability within species, it is recommended that the user conduct limited field tests before broad scale use.

COMPATIBILITY

Use caution if combined with highly alkaline materials such as lime, sulfur or dormant spray oils. Spraying concurrently with basic copper should be avoided. However, if it is to be tank mixed, a simple jar test should be conducted to test for compatibility. Mixing with certain other compounds could result in an incompatible mix.

APPLICATION INFORMATION

Formula 1 SW may be applied to crops by all application methods including conventional sprays, drip irrigation, soil application and aerial application. When applied by concentrate or dilute ground sprayers, use with a minimum of 15 gallons of water per acre. When used in aerial applications, use at least 5 gallons of water per acre. Control drift of this or other agricultural sprays. .If nitrogen is needed, add ten pounds of low biuret urea to 100 gallons of water.

Limitations, Restrictions, and Exceptions

RATES AND TIMING OF APPLICATION

Citrus and Avocados, Tree Fruit and Nut Crops: Apply 2 to 4 pints per acre per

application, beginning at pre-bloom and additional applications at two to four week intervals or as needed to meet nutritional requirements.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Drip Irrigation](#)

[Soil application](#)

Timings

[Beginning at pre-bloom.](#)