

# **RICE**

## General Information

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For use only in commercial seed treatment equipment. Not for use in hopper box, planter box, slurry box, or other on-farm seed treatment applications.

### SEED BAG TAG

The Federal Seed Act requires that the container of seed treated with EverGol Xtend C must be labeled with the following statements:

- This seed has been treated with EverGol Xtend C which contains Trifloxystrobin and Penflufen.
- Do not use treated seed for food, feed, or oil production. Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.
- User is responsible for ensuring that the seed bag meets all the requirements under the Federal Seed Act.

In addition, the US Environmental Protection Agency requires the following statements on the container of seed treated with EverGol Xtend C:

- Store away from feeds and foodstuffs.
- Wear long sleeved shirt, long pants, shoes, socks, and chemical resistant gloves when handling treated seed.
- Crops on this label and other crops for which tolerances exist for trifloxystrobin and penflufen may be replanted at any time. All other crops may be planted after 30 days.
- After the seeds have been planted, do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. Exception: Once the seeds are planted in soil or other planting media, the Worker Protection Standard allows worker to enter the treated area without restriction if there will be no worker contact with the treated seeds in the soil or planting media.
- Dispose of all excess treated seed. Left over treated seed may be double sown

around the headland or buried away from water sources in accordance with local requirements. Do not contaminate water bodies when disposing of planting equipment washwaters.

- Dispose of seed packaging in accordance with local requirements.
- Use planting equipment that will plant treated seed into the soil to a minimum depth of 0.5 inch.
- This compound is toxic to birds and mammals. Treated seed exposed on soil surface may be hazardous to birds and mammals. Cover or collect seeds spilled during loading.
- Do not plant treated rice seed directly into flooded field. Do not soak treated rice seed.
- For alfalfa: Do not exceed 0.00225 lb. penflufen /A (1.02 g ai/A) per crop season. The maximum number of applications per year is 3.
- For beans and peas (excluding soybeans): Do not exceed 0.0384 lb. penflufen /A (17.5 g ai/A) per crop season. The maximum number of applications per year is 3.
- For soybeans: Do not exceed 0.0384 lb. penflufen /A (17.5 g ai/A) per crop season. The maximum number of applications per year is 3.
- For Barley, Triticale, and Wheat (based on wheat seeding rate): Do not exceed 0.0078 lb. penflufen /A (3.5 g ai/A) per crop season. The maximum number of applications per year is 3.
- For Cotton (based on cotton seeding rate): Do not exceed 0.0019 lb. penflufen /A (0.86 g ai/A) per crop season. The maximum number of applications per year is 3.
- For Corn (field, sweet, popcorn) and sorghum (based on corn or sorghum seeding rate): Do not exceed 0.0033 lb. penflufen /A (1.5 g ai/A) per crop season. The maximum number of applications per year is 3.
- For rice (dry seeded): Do not exceed 0.006 lb. Penflufen /A (2.7 g ai/A) per crop season. The maximum number of applications per year is 3.
- This product has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow. Therefore contain any product spills or equipment leaks and dispose of wastes according to the disposal instructions on this label.

## INFORMATION

EverGol Xtend C is a fungicide seed dressing for protection against listed soilborne, seedborne, and early season post-emergence diseases of listed crop plants.

Limitations, Restrictions, and Exceptions

- Seed-borne false smut (suppression)
- Seed-borne kernel smut (suppression)

NOTE: To deliver 0.0010 – 0.0021 mg ai per seed, apply 0.011 – 0.023 fl oz of the product per 100,000 Rice seeds. (Based on an average of 21,850 Rice seeds per lb)

## DIRECTIONS

Apply using commercial slurry or mist-type seed treatment equipment. Uniform application of seed is necessary to ensure seed safety and best disease protection. Dilute product with sufficient water to ensure complete seed coverage.

Use high quality, properly cleaned seed.

All seed must be adequately dyed with a suitable colorant to prevent its accidental use as food for man or feed for animals. Refer to 21CFR, part 2.25. Any colorant added to treated seed must be cleared for use under 40CFR, part 153.155.

## Method

### [Seed Treatment](#)

### Restricted Entry Interval

12 hours

Exception: Once the seeds are planted in soil or other planting media, the Worker Protection Standard allows worker to enter the treated area without restriction if there will be no worker contact with the treated seeds in the soil or planting media.

## Timings

[N.A.](#)