

GRASS CUT FOR HAY - SUPPLEMENTAL LABEL

General Information

SPRAY DRIFT MANAGEMENT

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

Apply only as a medium or coarser spray (ASAE standard 572) of a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Apply only when the wind speed is 2 to 10 mph at the application site.

Additional Requirements for Aerial Applications

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter. Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy. When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

Do not make applications into temperature inversions.

Additional Requirements for Ground Boom Applications.

Do not apply with a nozzle height greater than 4 feet above the crop canopy.

SELECTIVE SPRAYING

NOTE: When using on grains and flax - do not forage or graze dairy and meat animals on treated areas within seven days of slaughter. Also, except for small underseeded grains, use at least 8 to 10 gallons of water per acre for ground application and at least 1 to 5 gallons of water per acre for aerial application.

Wheat, barley and oats treated with MCPA may be replanted with any crop specified on an MCPA label or any crop for which a residue tolerance exists for MCPA. For crops not listed on an MCPA label, or on crops for which no residue tolerances for

MCPA have been established, a 60-day plantback interval must be observed.

Limitations, Restrictions, and Exceptions

GRASS CUT FOR HAY

Grass Cut for Hay: Use 3/4 to 2 pints per acre in sufficient water (2 to 100 gallons of mixture) in airplane or ground spray application to give thorough coverage. Use higher rate for White top, Canada thistle and other less easy-to-kill weeds; spray perennials in early bud to full bloom stage and regrowth in Fall. Other weeds in Spring or Fall.

Precautions and Restrictions

When using on Grass Cut for Hay there is a (1) 7 day pre-grazing interval for dairy cattle; (2) 3 day pre-slaughter interval for meat animals.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Restricted Entry Interval

12 hours

Timings

[Spot treatment](#)

[Spot treatment](#)