

AGRICULTURE CROPS - FOLIAR SPRAY RECOMMENDATIONS

General Information

FORMULATED AS AN AID FOR HELPING PREVENT CORK SPOT, BITTER PIT, AND OTHER CALCIUM DEFICIENCY DISORDERS IN FRUIT TREES, FIELD AND VEGETABLE CROPS.

GENERAL INFORMATION:

Quick Cal is most effective when plants are growing rapidly and least effective when dormant or suffering from stress.

On plants with waxy leaf surfaces, use a nonionic wetting agent at recommended rates. Do not use while

temperatures are exceeding 90°F or when plants are under stress conditions.

Determine compatibility with other

fertilizers and/or pesticides by use of a jar test before tank mixing.

DIRECTIONS FOR USE:

Pour recommended amount of Quick Cal into partly filled spray tank. Add balance of water to fill tank. Keep agitated while filling and spraying. Flush all spray equipment after use.

Apply the product with non-corrosive spray equipment containing parts made of stainless steel, P.V.C., plastic, or fiberglass.

Limitations, Restrictions, and Exceptions

Foliar spray recommendations: Because of the nature of calcium deficiencies, Quick Cal should be applied in multiple

applications at 10-14 day intervals during the rapid growing and/or fruiting stage.

For best results, spray when the crop is

in an active growing state: after irrigation or natural rainfall. Spray early in the morning or late afternoon, Midday sprays

may not be effective because of excessive moisture evaporation.

Method

[Foliar spray](#)

Timings

Early in the morning or late afternoon

Early in the morning or late afternoon

Early in the morning or late afternoon