

ESTABLISHED PASTURES AND RANGELANDS

General Information

SPRAY DRIFT MANAGEMENT

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates. 2,4-D esters may volatilize during conditions of low humidity and high temperatures. Do not apply during conditions of low humidity and high temperatures.

For aerial application:

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter. Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety. This requirement does not apply to forestry or rights-of-way applications.

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this by adjusting the path of the aircraft upwind.

For ground boom application:

Do not apply with a nozzle height greater than 4 feet above the crop canopy.

GENERAL INFORMATION

This product is a low volatile ester especially prepared for use on crops and weeds where a susceptible crop in the near vicinity may be injured by a more volatile product. It is recommended for control of numerous broadleaf weeds and certain 2,4-D susceptible woody plants without injury to most established grasses. In cropland, 2,4-D is more effective than amines for controlling hard-to-kill weeds such as Bindweed, Curly dock, Smartweeds, Tansy ragwort, Thistle, Wild garlic, and Wild onions. For best results, apply this product as a water or oil spray during warm weather when young succulent weeds or brush are actively growing. Application under drought conditions often will give poor results. The lower recommended rates will be satisfactory on susceptible, annual weeds. For perennial weeds and conditions such as the very dry areas of the Western States where control is difficult, the higher recommended rates should be used. Deep-rooted perennial weeds such as Canada thistle and Field bindweed and many woody plants usually require repeated applications for maximum control.

Unless otherwise recommended, suggested application rates may be 1 to 10 gallons of total spray by air or 5 to 25 gallons by ground application equipment. If band treatment is used, base the dosage rate on the actual area to be sprayed. Although water quantities may vary due to different types of application equipment, sufficient water must be used to provide for complete and uniform coverage. In all cases, use the same recommended amount of 2,4-D per acre. When product is used for weed control in crops, the growth stage of the crop must be considered. For crop uses, do not mix with oil or other adjuvants unless specifically recommended on label. To do so may reduce herbicide's selectivity and could result in crop damage. If you are not prepared to accept some degree of crop injury, do not use this product.

Crop varieties vary in response to 2,4-D and some are easily injured. Apply this product to varieties known to be tolerant to 2,4-D. If you are uncertain concerning tolerant varieties or local use situations that may affect crop tolerance to 2,4-D, consult your seed company, State Agricultural Extension Service or qualified crop consultant for advice.

Aerial applications should be used only when there is no danger of drift to

susceptible crops. Many states have regulations concerning aerial application of 2,4-D formulations. Consult local regulatory authorities before making applications. Although this product is a low volatile formulation, at temperatures above 90°F vapors may damage susceptible crops growing nearby.

TO PREPARE THE SPRAY: (1) Fill the spray tank about half full with water, then add the required amount of this product with agitation, and finally the rest of the water.

NOTE: This product in water forms an emulsion which tends to separate unless the mixture is kept agitated. Continue agitation during application until spray tank is empty. (2) If oil is added, first mix this product and the oil and then add this mixture to the water. However, with adequate agitation, the oil can be added after the product is mixed in water. (3) If straight oil is used, a solution is formed and separation does not occur. Do not allow any water to get into the oil-herbicide mixture to avoid formation of an invert emulsion.

SELECTIVE WEEDING IN CROPS

USE IN LIQUID NITROGEN FERTILIZER: This product may be combined with liquid nitrogen fertilizer suitable for foliage application on corn, grass, pastures, or small grains in one operation. Use product according to directions on this label for those crops. Use liquid nitrogen fertilizer at rates recommended by supplier or Extension Service Specialist. Mix the product and fertilizer according to the following instructions: Fill the spray tank approximately half full with the liquid nitrogen fertilizer. Add the product while agitating the tank. Add the remainder of the fertilizer while continuing to agitate. Apply immediately, maintaining agitation during application until tank is empty. Do not apply during cold (near freezing) weather. Spray mixture must be used immediately and may not be stored. Do not allow mixture to stand overnight.

NOTE: If good, continuous agitation is not maintained, separation of the spray mixture and/or clogging of the nozzles is likely to occur. Fertilizers can increase foliage contact burn of herbicides. Reducing the fertilizer rate and concentration will reduce the hazard of leaf burn.

Limitations, Restrictions, and Exceptions

ESTABLISHED PASTURES AND RANGELANDS: The rates of application per acre per application per site. Use 2/3 to 2-2/3 pints of product in sufficient water to give good

coverage to one acre depending on type of weeds and stage of growth. Use only on established stands of perennial grasses. Do not use on alfalfa, bentgrass, clover, or other legumes. Do not use on newly seeded areas until grass is well established. Do not use from early boot to milk stage when grass seed production is desired.

Use Precautions for Pasture and Rangeland

Do not cut forage for hay within 7 days of application.

Postemergence:

For susceptible annual and biennial broadleaf weeds: Use 1.45 pints of product /acre per application.

For moderately susceptible biennial and perennial broadleaf weeds: Use 1.45 to 2.9 pints of product /acre per application.

For difficult to control weeds and woody plants: Use 2.9 pints of product /acre per application.

Spot treatment: Use 2.9 pints of product /acre.

Maximum of two applications per year.

Maximum of 5.8 pints of product /acre per year.

Minimum of 30 days between applications.

If grass is to be cut for hay, Agricultural Use Requirements for the Worker Protection Standard are applicable.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Restricted Entry Interval

12 hours

Timings

[N.A.](#)