

FIELD EQUIPMENT DISINFECTION (AGAINST CITRUS CANKER)

General Information

StorOx 2.0 works best when diluted with water with minimal levels of organic or inorganic materials, and with water having a neutral pH. Thoroughly rinse out tank with water before mixing concentrate. StorOx 2.0 will readily mix with clean, neutral water and does not require agitation.

StorOx 2.0 concentrate should not be combined or mixed with any other pesticide concentrates.

Limitations, Restrictions, and Exceptions

FIELD EQUIPMENT DISINFECTION (AGAINST CITRUS CANKER)

StorOx 2.0 may be used to disinfect harvest equipment such as pickers, trailers, trucks (including truck body parts and tires), bins, packing crates, ladders, power tools, hand tools, gloves, rubber boots, pruning shears or other equipment that may transfer *Xanthomonas campestris* (axonopodis) pathovars citrumelo (citrus canker surrogate). This product can also be used to disinfect surfaces contaminated with *P. aeruginosa*, *S. enterica* and *S. aureus*.

1. Remove gross contamination with a cleaner or other suitable detergent and rinse with water.
2. For *Xanthomonas campestris* (axonopodis) pathovars citrumelo (citrus canker surrogate), use StorOx 2.0 at a dilution rate of 1:400–1:800 as a general coarse spray. For *P. aeruginosa*, *S. enterica* and *S. aureus*, use StorOx 2.0 at a dilution rate of 1:100 (1.3 fl. oz./gal) as a general coarse spray.
3. Allow StorOx 2.0 to contact surface for ten (10) minutes.
4. Allow to air dry, do not rinse.

Method

[Spray](#)

Timings

[N.A.](#)