

POST HARVEST APPLICATIONS - APPLICATION TO WATER USED FOR SPRAYING OF FRUITS AND VEGETABLES

General Information

StorOx 2.0 works best when diluted with water with minimal levels of organic or inorganic materials, and with water having a neutral pH. Thoroughly rinse out tank with water before mixing concentrate. StorOx 2.0 will readily mix with clean, neutral water and does not require agitation.

StorOx 2.0 concentrate should not be combined or mixed with any other pesticide concentrates.

Limitations, Restrictions, and Exceptions

POST HARVEST APPLICATIONS

Use StorOx 2.0 to treat bacterial and fungal diseases in post-harvest waters, systems, equipment, structures and on agricultural commodities.

FOR TREATMENT OF FRUIT AND VEGETABLE SURFACES AND WASH OR PROCESS WATERS

Use StorOx 2.0 to control the growth of post-harvest spoilage and decay-causing bacterial and fungal plant pathogens on fruits and vegetables. Product can be added directly to the wash water or spray water used for cleaning of fruits and vegetables.

APPLICATION TO WATER USED FOR SPRAYING OF FRUITS AND VEGETABLES

Mix StorOx 2.0 with water used for spray wash water at a dilution rate of 0.35–0.58 fl. oz. per gallon of water. Spray until even and sufficient distribution is achieved on the produce surface and a contact time of at least 10 seconds with the applied solution. Allow produce surface to air dry to ensure no residual moisture prior to packing and/or storage.

Method

[Spray](#)

Timings

[N.A.](#)