

FOLIAR APPLICATION - FIELD AND ROW CROPS

General Information

DIRECTIONS FOR USE

Actagro Micro Booster can be applied to most vegetable and cucurbit crops, field and row crops, deciduous fruit and nut trees, grapes, berries, olives, avocados and citrus. Micro Booster is designed for soil application and for foliar use. Apply directly to soils or inject through irrigation systems where water quality is satisfactory to maintain solubility. Foliar applications should be as a dilute spray. Application rates vary by crop or situation. Mixing with irrigation water for field storage is generally not recommended. Products blended with irrigation water should be used immediately and not stored for prolonged periods.

GENERAL INFORMATION

Micro Booster will be influenced by various soil and environmental conditions and should be applied under conditions of optimum plant growth. Optimum conditions are defined as adequate humidity, moderate temperatures and adequate soil moisture. Optimum conditions for most crops generally occur during early morning or late evening hours.

CAUTION: If any other chemicals are to be used in the spray mixture be sure to check compatibility. Foliar application of Micro Booster close to harvest may leave a brown residue on the harvestable portion of the crop. Rainfall, sprinkler irrigation or high humidity should alleviate this condition.

Limitations, Restrictions, and Exceptions

FOLIAR APPLICATION

FIELD & ROW CROPS: Apply in a minimum of 10 gallons of water per acre. May be applied through center pivot irrigation systems. Repeat applications may be beneficial.

Method

[Foliar application](#)

Timings

N.A.