

# **PEACH PREPLANT ROOT DIP (CALIFORNIA)**

## General Information

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Read all precautions and directions for use before using. Use only for claims listed and only as specified on this label.

In order that pesticide residues on food and forage crops will not exceed federal tolerances, use only at recommended rates and intervals and do not apply closer to harvest than specified. Do not apply or allow to drift to adjoining food, fiber, or pasture crops. Drift of CAPTAN 80WDG onto sensitive crops (e.g., D'Anjou Pears) can cause severe phytotoxicity and crop loss.

### SPRAY DRIFT LABELING

Do not allow this product to drift.

#### Foliar Spray Drift Management

Avoiding spray drift from foliar applications is the responsibility of the applicator. Similar to aerial spray drift, the interaction of many equipment- and-weather-related factors determine the potential for spray drift from foliar applications. To protect water resources, the applicator and the grower are responsible for considering all these factors when making decisions.

#### Aerial Spray Drift Management

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed  $\frac{3}{4}$  the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory Information.

#### Aerial Drift Reduction Advisory

This section is advisory in nature and does not supersede the mandatory label requirements.

**COMPATIBILITY AND PLANT SAFETY:** CAPTAN 80WDG can be combined safely and effectively at recommended dosage rates with most commonly used fungicides and insecticides with the exception of oil and strongly alkaline materials. Alkaline materials such as spray lime, lime-sulfur, and Bordeaux mixture will reduce the fungicidal activity of CAPTAN 80WDG. Do not apply CAPTAN 80WDG in combination with or immediately before or closely following oil sprays. Do not allow oil sprays on adjacent crops to drift onto crops which have been or will shortly be treated with CAPTAN 80WDG. The time factor governing the safe interval between CAPTAN 80WDG and oil sprays varies due to general climatic conditions, therefore, consult local agricultural spray programs and authorities to determine the proper timing. The use of spreaders which cause excessive wetting is not advised. Combinations with solvent formulations of organic phosphates should not be used. Combinations of CAPTAN 80WDG and sulfur should not be used on crops sensitive to sulfur. Used at high rates or in drenching sprays, CAPTAN 80WDG may cause a necrotic spotting of tender, immature leaves of certain varieties of apples, peaches, plums, and cherries. This type of injury is most likely to occur in the early cover sprays during long periods of warm, cloudy, humid weather. To avoid the hazard of leaf spotting under such conditions, use CAPTAN 80WDG and other spray materials at lowest recommended rates and avoid drenching trees.

Applications can be made by aircraft or ground power equipment (including concentrate and semi-concentrate equipment). Pour recommended amount of this material into nearly filled spray tank. Add balance of water. Maintain agitation during filling and spraying operations. Do not allow mixture to stand. Do not combine with emulsifiable liquids or wettable powders unless previous experience has proven them to be physically compatible and safe to plants. (Read compatibility and plant safety information.)

For aerial or concentrate spray applications, apply the same amount of CAPTAN 80WDG per acre as would normally be applied for dilute spray applications. Apply aerial or concentrate sprays in sufficient water for coverage.

#### CHEMIGATION STATEMENT

Do not apply this product through any type of irrigation system.

#### GENERAL USE PRECAUTIONS

Except as specified, begin applications before or at first sign of disease and repeat as needed to maintain control but observe use limitations. Unless otherwise specified, application can be made on the day of harvest. Maximum application is for a crop cycle. Crop cycle is defined as prebloom through postharvest. Apply the high rate and/or spray at shorter intervals when climatic conditions most favor disease(s). Apply the low rate and/or spray at larger intervals when climatic conditions least favor disease(s). If you are unaware of the climatic conditions favorable for disease(s) claimed for the specific use sites, you must consult with your State Agricultural Extension Service to learn of these conditions.

**IMPORTANT:** Read label carefully. Although most of the directions on this label may be followed nationwide, a few are limited to either the eastern or western U.S. Follow those directions for your growing area where specified.

If applying as a directed/banded spray; use band rate of CAPTAN 80WDG according to the following formula given in the label.

#### Limitations, Restrictions, and Exceptions

**PEACH PREPLANT ROOT DIP (California):** Preventative preplant dip treatment for crown gall; Use 2½ pounds CAPTAN 80WDG plus 3.2 pints sodium hypochlorite (5.25% household bleach) per 100 gallons of water. Wash nursery trees to remove soil from roots. Cut off all dormant buds and suckers in crown area and prune root

system if necessary. Submerge the entire dormant tree for 5 minutes. Recharge dip during operation at a rate of 3.2 pints diluted sodium hypochlorite per 100 gallons of water.

Method

[Root Dip](#)

Timings

[Preplant](#)