

IN AND AROUND BUILDINGS AND STRUCTURES - FOR MICE

General Information

ENDANGERED SPECIES CONSIDERATION

Notice: The use of this product may pose a hazard to Federally designated endangered/ threatened species. It is a Federal offense to use any pesticide in a manner that results in the death of a member of an endangered species. Consult the nearest U.S. Fish and Wildlife Service regional office or the appropriate State Agency for current information on habitats occupied by endangered species.

Whooping Crane (*Grus americana*). Do not use this product in habitats occupied or occasionally visited by whooping cranes during the period from 30 days before the expected arrival of cranes to 30 days after the time of their usual departure.

Black-footed Ferret (*Mustela nigripes*). Do not use this product within 7 kilometers (4.34 miles) of any prairie dog town to limit risks to the black-footed ferret from exposure to Zinc Phosphide or destruction of its prey base, unless the colony is an isolated black-tailed prairie dog town less than 80 acres in size or an isolated white-tailed prairie dog town less than 200 acres in size, or unless the town had been appropriately surveyed, using methods acceptable to the U.S. Fish and Wildlife Service, and found by the FWS not to be suitable site for ferret reintroductions.

Gray Wolf (*Canis lupus*) and Grizzly Bear (*Ursus arctos horribilis*). Unless the local U.S. Fish and Wildlife Service office has determined that there are no gray wolves or grizzly bears in the general vicinity of bait applications in Montana and Wyoming, do not apply this product outdoors within occupied habitat of these species.

Attwater's Greater Prairie Chicken (*Tympanuchus cupido attwateri*). Do not use this product in the present occupied range of the Attwater's Greater Prairie Chicken in the following counties: Aransas, Austin, Brazoria, Colorado, DeWitt, Ft. Bend, Galveston, Goliad, Hays, Refugio, Victoria, Waller, and Wharton (Texas).

Yellow-Shouldered Blackbird (*Agelaius xanthomus*) and Puerto Rican Plain Pigeon (*Columba inornata wetmorei*). This product must not be used in the following areas in Puerto Rico except in tamper-proof bait boxes:

(1) within 6.3 miles (10 kilometers) of Central Aquirre, Lago Cidra, Ceiba, San

German

(2) within 9.4 miles (15 kilometers) of La Esperanza, south of Highway 2 from city of Mayaguez to the city of Ponce and all Mona Island

Utah Prairie Dog (*Cynomys parvidens*). Do not use this product in critical habitat of the Utah Prairie Dog (Utah).

Salt Marsh Harvest Mouse (*Reithrodontomys raviventris*). Do not use this product in critical habitat within 0.5 miles of salt marsh vegetation and/or brackish water wetlands which are located: 1) near or adjacent to San Pablo Bay and San Francisco Bay, or 2) in the Sacramento River below or adjacent to confluence of the Sacramento River and the San Joaquin River (California)

Morro Bay Kangaroo Rat (*Dipodomys heermanni morroensis*). Do not use this product in critical habitat within 2.5 miles of Baywood Park which is located on Morro Bay (California)

Giant Kangaroo Rat (*Dipodomys ingrens*). Follow the Interim Measures for protecting endangered species in the California Endangered Species Bulletin for the following California counties: Fresno, Kern, Kings, Merced, Monterey, San Benito, San Luis Obispo, Santa Barbara and Tulare (California)

Tipton Kangaroo Rat (*Dipodomys nitratoides nitratoides*). Follow the Interim Measures for protecting endangered species in the California Endangered Species Bulletin for the following California counties: Kern, Kings and Tulare (California)

Fresno Kangaroo Rat (*Dipodomys nitratoides exilis*). Follow the Interim Measures for protecting endangered species in the California Endangered Species Bulletin for the following California counties: Fresno, Kern, Kings, Madera and Merced (California)

Stephan's Kangaroo Rat (*Dipodomys stephens*). Follow the Interim Measures for protecting endangered species in the California Endangered Species Bulletin for the following California counties: San Bernardino, San Diego and Riverside (California)

San Bernardino Kangaroo Rat (*Dipodomys merriami parvus*). Follow the Interim Measures for protecting endangered species in the California Endangered Species Bulletin for the following California counties: San Bernardino and Riverside (California)

Northern Idaho Ground Squirrel (*Spermophilus brunneus brunneus*). Do not use this product within 0.5 miles in critical habitat of the Northern Idaho Ground Squirrel in Adams and Valley counties, Idaho.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions. Use only for the sites, pests, and application methods described on this label.

Limitations, Restrictions, and Exceptions

ENDANGERED SPECIES CONSIDERATION

Notice: The use of this product may pose a hazard to Federally designated endangered/ threatened species. It is a Federal offense to use any pesticide in a manner that results in the death of a member of an endangered species. Consult the nearest U.S. Fish and Wildlife Service regional office or the appropriate State Agency for current information on habitats occupied by endangered species.

Whooping Crane (*Grus americana*). Do not use this product in habitats occupied or occasionally visited by whooping cranes during the period from 30 days before the expected arrival of cranes to 30 days after the time of their usual departure.

Black-footed Ferret (*Mustela nigripes*). Do not use this product within 7 kilometers (4.34 miles) of any prairie dog town to limit risks to the black-footed ferret from exposure to Zinc Phosphide or destruction of its prey base, unless the colony is an isolated black-tailed prairie dog town less than 80 acres in size or an isolated white-tailed prairie dog town less than 200 acres in size, or unless the town had been appropriately surveyed, using methods acceptable to the U.S. Fish and Wildlife Service, and found by the FWS not to be suitable site for ferret reintroductions.

Gray Wolf (*Canis lupus*) and Grizzly Bear (*Ursus actos horribilus*). Unless the local U.S. Fish and Wildlife Service office has determined that there are no gray wolves or grizzly bears in the general vicinity of bait applications in Montana and Wyoming, do not apply this product outdoors within occupied habitat of these species.

Attwater's Greater Prairie Chicken (*Tympanuchus cupido attwateri*). Do not use this

product in the present occupied range of the Attwater's Greater Prairie Chicken in the following counties: Aransas, Austin, Brazoria, Colorado, DeWitt, Ft. Bend, Galveston, Goliad, Hays, Refugio, Victoria, Waller, and Wharton (Texas).

Yellow-Shouldered Blackbird (*Agelaius xanthomus*) and Puerto Rican Plain Pigeon (*Columba inornata wetmorei*). This product must not be used in the following areas in Puerto Rico except in tamper-proof bait boxes:

(1) within 6.3 miles (10 kilometers) of Central Aquirre, Lago Cidra, Ceiba, San German

(2) within 9.4 miles (15 kilometers) of La Esperanza, south of Highway 2 from city of Mayaguez to the city of Ponce and all Mona Island

Utah Prairie Dog (*Cynomys parvidens*). Do not use this product in critical habitat of the Utah Prairie Dog (Utah).

Salt Marsh Harvest Mouse (*Reithrodontomys raviventris*). Do not use this product in critical habitat within 0.5 miles of salt marsh vegetation and/or brackish water wetlands which are located: 1) near or adjacent to San Pablo Bay and San Francisco Bay, or 2) in the Sacramento River below or adjacent to confluence of the Sacramento River and the San Joaquin River (California)

Morro Bay Kangaroo Rat (*Dipodomys heermanni morroensis*). Do not use this product in critical habitat within 2.5 miles of Baywood Park which is located on Morro Bay (California)

Giant Kangaroo Rat (*Dipodomys ingens*). Follow the Interim Measures for protecting endangered species in the California Endangered Species Bulletin for the following California counties: Fresno, Kern, Kings, Merced, Monterey, San Benito, San Luis Obispo, Santa Barbara and Tulare (California)

Tipton Kangaroo Rat (*Dipodomys nitratoide nitratoide*). Follow the Interim Measures for protecting endangered species in the California Endangered Species Bulletin for the following California counties: Kern, Kings and Tulare (California)

Fresno Kangaroo Rat (*Dipodomys nitratoide exilis*). Follow the Interim Measures for protecting endangered species in the California Endangered Species Bulletin for the following California counties: Fresno, Kern, Kings, Madera and Merced (California)

Stephan's Kangaroo Rat (*Dipodomys stephens*). Follow the Interim Measures for protecting endangered species in the California Endangered Species Bulletin for the following California counties: San Bernardino, San Diego and Riverside (California)

San Bernardino Kangaroo Rat (*Dipodomys merriami parvus*). Follow the Interim Measures for protecting endangered species in the California Endangered Species Bulletin for the following California counties: San Bernardino and Riverside (California)

Northern Idaho Ground Squirrel (*Spermophilus brunneus brunneus*). Do not use this product within 0.5 miles in critical habitat of the Northern Idaho Ground Squirrel in Adams and Valley counties, Idaho.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions. Use only for the sites, pests, and application methods described on this label.

USES ASSOCIATED WITH BUILDINGS AND STRUCTURES TO CONTROL NORWAY RATS, ROOF RATS, AND HOUSE MICE

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from bait stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooved livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.

3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

Note: Bait stations are mandatory for outdoor, above-ground use. Tamper-resistant bait stations must be used if children, pests, non-target mammals, or birds may access the bait.

USE RESTRICTIONS:

IN AND AROUND BUILDINGS AND STRUCTURES

This product may only be used to control Norway rats, roof rats and house mice in and within 100 feet of man-made structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to harboring or attracting rodent infestations. Examples of such structures include homes and other permanent or temporary residences, food processing facilities, industrial and commercial buildings, trash receptacles, agricultural and public buildings, transport vehicles (ships, trains, aircraft), docks and port of terminal buildings and related structures around and associated with these sites. Fence and perimeter baiting, beyond 100 feet from a structure as defined above, is prohibited. This product must not be applied directly to food or feed crops. Do not broadcast bait.

Inside Burrows

This product may be applied to active rodent burrows to control Norway rats, and roof rats within or beyond 100 feet of buildings and man-made structures, provided that infestations of these rodents have been confirmed. Because Norway/roof rat infestations may occur in areas farther than 100 feet from buildings and man-made structures when the rodents have ample supplies of food and cover, efforts should be made to remove food trash, garbage, clutter and debris. Bait must be placed not less than 6 inches into active Norway/roof rat burrows. Do not broadcast bait.

SELECTION OF TREATMENT AREAS: After removing as much food, water and harborage as possible, determine dry, acid-free areas where rats and mice will most likely find and consume the bait. Generally these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rats and mice or their signs have been observed.

APPLICATION DIRECTIONS FOR MICE: Apply 1 teaspoon of bait per placement at 8 to 12 foot intervals. For outdoor use, place bait in tamper-resistant bait stations. Maintain an uninterrupted supply of fresh bait until all signs of feeding have

stopped. Do not treat the same area at less than 30 day intervals.

FOLLOW-UP: Using waterproof gloves, collect and properly dispose of all dead animals and excess bait properly in accordance with "Pesticide Disposal" instructions. Use detergent and hot water to wash spoons for application into burrows. Do not use spoon for mixing, holding or transferring food or feed. To discourage reinfestation, limit sources of rodent food, water, and harborage as much as possible.

USES ASSOCIATED WITH NON-BUILDING AGRICULTURAL SITES Use Restrictions for all Non-Structural Agricultural Sites

- Do not apply this product in a way that will contact workers or other persons, either directly or through drift.
- Only protected handlers may be in the area during application.
- Keep all other persons out of the treated area during this application.
- Do not apply on roads, near residential areas, or over water.
- Do not broadcast over crops unless use directions specifically permit aerial application.
- Apply bait on warm clear days.

USES ASSOCIATED WITH NON-BUILDING AGRICULTURAL SITES Use Restrictions for all Non-Structural Agricultural Sites

- Do not apply this product in a way that will contact workers or other persons, either directly or through drift.
- Only protected handlers may be in the area during application.
- Keep all other persons out of the treated area during this application.
- Do not apply on roads, near residential areas, or over water.
- Do not broadcast over crops unless use directions specifically permit aerial application.
- Apply bait on warm clear days.

Method

[Bait](#)

Timings

[N.A.](#)