

ADDITION OF BERMUDAGRASS MITE AS A PEST CONTROLLED ON GOLF COURSE GREENS, TEES, AND FAIRWAYS - AL, AR, AZ, CA, ETC.

General Information

PRODUCT INFORMATION

- Divanem is a suspension concentrate containing abamectin which, when applied according to directions for use, will control turf-parasitic nematodes such as sting (*Belonolaimus longicaudatus*), lance (*Hoplolaimus galeatus*), root-knot (*Meloidogyne* spp.), and ring (*Criconemoides* spp.) on golf courses and professional and collegiate sports fields.
- Divanem should be applied as a seasonal program.
- Apply Divanem as an early curative treatment (after appropriate nematode extraction, identification, and counts).
- Multiple applications may be required before improvements in turf quality are observed.
- Maximum annual rate must not exceed 0.27 lb abamectin/calendar year or 50 fl oz Divanem/A/calendar year.

APPLICATION DIRECTIONS

Application Equipment

For best results, use spray nozzles that produce extremely coarse droplets [such as Turbo Floodjet (TF5) or TurfJet (TTJ10)].

RESTRICTIONS AND PRECAUTIONS

Use Restrictions

- DO NOT apply to golf course roughs.
- DO NOT apply to residential turf, sports fields (other than professional or collegiate), or commercial turf.
- DO NOT use aerial application.
- Chemigation: DO NOT apply this product through any type of irrigation system.
- DO NOT apply through an ultra-low volume spray system.
- DO NOT apply to turf under heat or moisture stress.
- DO NOT GRAZE TREATED TURF OR FEED TURF CLIPPINGS FROM ANY TREATED

AREA TO POULTRY OR LIVESTOCK.

Limitations, Restrictions, and Exceptions

ADDITION OF BERMUDAGRASS MITE AS A PEST CONTROLLED ON GOLF COURSE GREENS, TEES, AND FAIRWAYS

AL, AR, AZ, CA, CO, CT, DC, DE, FL, GA, HI, IA, ID, IL, IN, KS, KY, LA, MA, MD, ME, MI, MN, MO, MS, MT, NC, ND, NE, NH, NJ, NM, NV, OH, OK, PA, RI, SC, SD, TN, UT, VA, VT, WA, WI, WV, WY

Directions for Use

For control of Bermudagrass mite (*Eriophyes* spp.) on golf course greens, tees, and fairways, apply Divanem at 3.125 to 6.25 fl oz/acre on a 14 day interval. Initiate applications prior to or in the early stages of symptom development. Tank mix with a surfactant or wetting agent and water in with 0.1 to 0.25 inch of irrigation immediately after application.

Follow all other directions for use, restrictions and precautions on the Divanem Section 3 label.

Method

[Spray](#)

Timings

[Prior to or in the early stages of symptom development.](#)