

HEAD AND STEM BRASSICA VEGETABLES - LOOPER ALFALFA, ETC.

General Information

GENERAL INSTRUCTIONS

Use low rate under light to moderate infestation. Higher rates should be used under heavy insect pressure. The rate of application is variable according to insect pressure, timing of spray and field scouting. Do not exceed maximum allowable rate.

Under hot and dry conditions, increase the spray volume and add appropriate spray adjuvant, but only to those crops which have previously demonstrated a tolerance to the adjuvant(s) under these conditions.

Preventive Use

For cutworm, armyworm, or stalk borer control, Mustang insecticide may be applied before, during, or after planting. For soil-incorporated applications, use higher rates for improved control. Do not exceed maximum allowable rate.

Refer the specified label of Maximum Seasonal Usage and PHI for all Crops.

Limitations, Restrictions, and Exceptions

HEAD AND STEM BRASSICA VEGETABLES

Apply in water as necessary for insect control using a minimum of 15 gallons of finished spray per acre with ground equipment and 5 gallons per acre by air.

Lower rates of Mustang should be used under light to moderate insect pressure.

Higher rates should be used to control heavy to extremely heavy insect populations.

In areas where arid climatic conditions persist, such as California and Arizona, higher than minimum recommended rates may be required.

- Do not make applications less than 7 days apart.

Aphids: Aphid control may be variable depending on species present and host-plant

relationships.

Whiteflies: Aids in control

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

1 day

Rates

[field_rates 0](#)

[field_rates 1](#)

•

Restricted Entry Interval

12 hours

Timings

[N.A.](#)