

WARM-SEASON TURFGRASSES - ANNUAL BLUEGRASS (POA ANNUA) CONTROL IN NON-OVERSEEDED TURF

General Information

GENERAL INFORMATION

Product Description: This product is a postemergence, systemic herbicide with limited soil residual activity. It provides postemergence control of many annual and perennial sedges, grass and broadleaf weeds on highly managed turf, sod farms and native grass sites. It is a selective herbicide that can be used over the top of many perennial warm-season turfgrasses. Refer to the appropriate sections of the label for approved turf species.

Use Sites: This product may be used for general weed control on highly managed turf, native grasses, landscaped areas and ornamental nurseries. This product may be applied to residential and commercial sites, including apartment complexes, athletic fields, cemeteries, golf course fairways, golf course roughs, golf course tees, and other golf course areas, hotel properties, nurseries, office complexes, parks, public areas, retail sites, storage facilities, school grounds, sod and turfgrass seed farms, and other highly managed turfgrass sites. This product is not for use on golf course putting greens.

When to Spray: Best results are obtained when target weeds are actively growing and not disturbed by mowing for at least 2 days before and 2 days after application.

Time to Symptoms: This product is absorbed by both the roots and the foliage of plants and rapidly inhibits the growth of susceptible weeds. Susceptible weed growth stops within 24 hours of treatment even though visual symptoms are slow to develop. Susceptible weeds usually show yellowing or browning within 2 to 3 weeks. Warm, moist conditions following application will accelerate herbicidal activity. Cold, dry conditions will delay herbicidal activity. Weeds stressed by drought are less susceptible to this product.

Rainfastness: Heavy rainfall or irrigation within 2 hours after application may wash this product off of the foliage and a repeat application may be required for adequate control.

Maximum Annual Use Rate: The combined total of all treatments must not exceed 2.66 ounces of this product per acre per year.

IMPORTANT : Avoid contact of this product with the roots or foliage of susceptible non-target vegetation as injury may occur. This includes areas where this product may be washed or moved into contact with roots of desirable vegetation.

Susceptible plants may be injured if seeded or transplanted into treated areas unless otherwise directed in the label.

Mixing for Hand-Held Sprayers

Hand-held sprayer applications should be made at a rate of 2 gallons of spray solution per 1000 square feet.

Using the measuring scoop provided in the product packaging, follow the instructions below to prepare the proper spray solution.

APPLICATION EQUIPMENT AND TECHNIQUES

Apply spray solutions of this product with properly maintained and calibrated equipment capable of delivering desired volumes.

Do not apply this product through any type of irrigation system.

Ground Broadcast Equipment

Apply this product uniformly with properly calibrated ground equipment in 10 to 50 gallons of water per acre. Select spray volumes that ensure thorough and uniform weed coverage. Use equipment that is capable of continuous agitation. Choose nozzles that provide optimum spray distribution and uniform coverage at a spray pressure appropriate for applying postemergence herbicides. Avoid streaking, skips, overlaps and spray drift during application.

Hand-Held Sprayers

Backpack sprayers, pump-up pressure sprayers, and other, similar types of hand-

held sprayers may be used to apply this product. Apply to foliage of vegetation to be controlled at a rate of 2 gallons of spray solution per 1000 square feet. Spray coverage should be uniform and complete. Do not spray to the point of runoff.

Refer in the label regarding tank mix information.

Limitations, Restrictions, and Exceptions

WARM-SEASON TURFGRASSES

- This product generally has been shown to be safe for use on the established warmseason turfgrasses listed in this section.
- Use of this product may result in temporary chlorosis, and may affect the growth pattern or delay green-up of the desirable turf. St. Augustinegrass and seashore paspalum may be more sensitive to this product than other grasses depending on environmental conditions, cultivar differences and other influential factors. For St. Augustinegrass and seashore paspalum, test this product on a small area prior to wide-scale use to determine if this product is suitable for your management and cultural practices.

Annual Bluegrass (*Poa annua*) Control in Non-Overseeded Turf

For selective control of annual bluegrass (*Poa annua*), apply this product at 1.25 to 2.0 ounces per acre. Use the higher rate of this product for control in areas of established, dense weed infestation.

Best results are obtained when weeds are in the early stage of growth and prior to tillering.

IN DORMANT BERMUDAGRASS ONLY, tank mixtures of this product with Roundup PROMAX, or Roundup QuikPRO herbicides may be used to increase the spectrum of vegetation controlled. Read and follow the label directions, precautionary statements and all other label information on Roundup PROMAX, or Roundup QuikPRO herbicides. Refer to the Roundup PROMAX, or Roundup QuikPRO product labels for approved application rates. Always apply tank mixtures according to the most restrictive precautionary statements of the products being used.

Method

[Broadcast/Foliar Ground](#)

Hand-Held Spray

Rates

field_rates 0

-

Restricted Entry Interval

12 hours

Timings

When weeds are in the early stage of growth and prior to tillering.