

CONSERVATION RESERVE PROGRAM - RENOVATION OF EXISTING PERENNIAL GRASSES

General Information

USE INFORMATION

ATRA-5 Herbicide will control most annual Broadleaf and Grass weeds in Corn, Sorghum, Sugarcane and certain other crops as specified on the label. This product may be applied before or after weeds emerge.

In each case where a range of rates is given, the lower rate should be used on Coarse textured soil and soil low in organic matter and the higher rate should be used on Fine textured soil and soil high in organic matter.

Since this product acts mainly through root absorption, its effectiveness depends on rainfall or irrigation to move it into the root zone. Should weeds develop, a shallow cultivation or rotary hoeing will generally result in better weed control.

This product is non-corrosive to equipment and metal surfaces, nonflammable and has low electrical conductivity. Care should be taken to avoid using this product near adjacent desirable plants or in greenhouses, or injury may occur.

To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, as crop injury may result.

Note: The Seller does not recommend the use of this product in combination with other herbicides or oils except as specifically described on the label or in literature published by the Seller.

FAILURE TO FOLLOW ALL USE PRECAUTIONS AND RESTRICTIONS ON THE LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY OR ILLEGAL RESIDUES.

RESISTANCE MANAGEMENT

This product is a Group 5 Herbicide.

Following many years of continuous use of this product and chemically related products, biotypes of some of the weeds listed on the label have been reported

which cannot be effectively controlled by this and related herbicides. Where this is known or suspected and weeds controlled by this product are expected to be present along with resistant biotypes, use this product in combinations or in sequence with other registered herbicides which are not triazines. If only resistant biotypes are expected to be present, use a registered non-triazine herbicide. Consult with your State Agricultural Extension Service for specific recommendations.

APPLICATION PROCEDURES

GROUND APPLICATION: Use conventional ground sprayers equipped with nozzles that provide accurate and uniform application. Be certain that nozzles are uniformly spaced and are the same size. Calibrate sprayer before use and recalibrate at the start of each season and when changing carriers. Unless otherwise specified, use a minimum of 10 gallons of spray mixture per acre for all pre-plant incorporated, preplant surface, pre-emergence and post-emergence applications (with or without oil or surfactant) with ground equipment. Use a pump with capacity to:

1. Maintain 35 to 40 psi at nozzles.
2. Provide sufficient agitation in tank to keep mixture in suspension.
3. Provide a minimum of 20% bypass at all times.

Use centrifugal pumps which provide propeller shear action for dispersing and mixing this product. The pump should provide a minimum of 10 gallons per minute per 100 gallon tank size circulated through a correctly positioned sparger tube or jets.

Use screens to protect the pump and to prevent nozzles from clogging.

Screens placed on the suction side of the pump should be 16-mesh or coarser. Do not place a screen in the recirculation line. Use 50-mesh or coarser screens between the pump and boom and where required, at the nozzles. Check nozzle manufacturer's recommendations.

AERIAL APPLICATION: Use aerial application only where broadcast applications are specified. Apply in a minimum of 1 quart of water for each quart of this product applied per acre. For post-emergence treatments on Corn and Sorghum, apply the rate in a minimum of 2 gallons of water per acre. Avoid applications under conditions where uniform coverage cannot be obtained or where excessive spray

drift may occur.

Avoid application to humans or animals. Flagmen and loaders should avoid inhalation of spray mist and prolonged contact with skin.

THIS PRODUCT IN WATER OR LIQUID FERTILIZER APPLICATIONS

Nitrogen solutions or complete liquid fertilizers may replace all or part of the water as a carrier for pre-emergence and pre-plant applications of this product on Corn and Sorghum. Check the compatibility of this product with liquid fertilizer and/or nitrogen solution as shown below before use. Do not apply in nitrogen solution or complete liquid fertilizer after Corn or Sorghum emerges, or crop injury may occur.

Refer in the label for tank mix information.

Limitations, Restrictions, and Exceptions

CONSERVATION RESERVE PROGRAM (CRP) (NE, OK, OR, TX) For control or suppression of the following weeds: Annual ragweed, Barnyardgrass, Black nightshade, Cheat, Cocklebur, Downy brome, Fall panicum, Field pennycress, Giant foxtail, Japanese brome, Kentucky bluegrass, Kochia, Lambsquarters, Little barley, Marehail, Pigweed, Prickly lettuce, Smooth brome, Sunflower and Yellow foxtail, refer to the directions for use, \"Notes\" and \"Use Precautions and Restrictions\" below:

Renovation of existing stands of the following Perennial range grasses planted on CRP acres: Bluegrama, Indiangrass, Little bluestem, Sand lovegrass, Side-oats grama and Western wheatgrass. Broadcast 0.8 to 1.6 pints of this product per acre in the Spring prior to weed emergence or in the Fall before the ground freezes and prior to weed emergence after these species have been established for at least one growing season for control or partial control of the weeds listed above. Use the lower rate for weeds controlled or suppressed easily. Use the higher rate on other weeds claimed in an earlier section of the label.

Aerial Application: Make applications at a maximum height of 10 feet above vegetation. Use low-drift nozzles at a maximum pressure of 40 psi. Restrict application to periods when wind speed does not exceed 10 mph to control drift. To assure that drift will not adversely affect adjacent sensitive non-target plants, apply this product by aircraft at a minimum upwind distance of 400 feet from sensitive

plants. Use 3 to 5 gallons per acre total water volume; use the higher water volume when a dense, heavy ground cover is present.

USE PRECAUTIONS AND RESTRICTIONS

1. Do not cut or feed Grass hay to livestock.
2. Do not graze treated areas.
3. Do not use seeds for bird food.
4. Do not dump or spill product or dispose of containers within reach of livestock.
5. Follow all applicable restrictions for the Conservation Reserve Program.
7. Do not apply more than 2 pounds a.i./A (3.2 pints/A) for any application.
8. Do not make more than one application per year. Slight discoloration of desirable grasses may occur following treatment. Injury may be enhanced when used on neutral or alkaline soils.

SEVERE DROUGHT CONDITIONS Do not graze forage or cut forage for hay. Under severe drought conditions, the Conservation Reserve Program allows grazing and making of hay from CRP acres, as so specified by the local ASCS (Agricultural Stabilization & Conservation Service) office. The label does not allow grazing or making of hay from CRP acres that have been treated with atrazine under any circumstances.

Method

[Broadcast/Foliar Air](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

12 hours

Exception: If the product is soil-injected or soil-incorporated, the WPS, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[Preemergence \(Weed\)](#)