

## **HEAD LETTUCE - APHID, ARMYWORM, EARWORM ETC.**

### General Information

#### GENERAL INSTRUCTIONS

Use low rate under light to moderate infestation. Higher rates should be used under heavy insect pressure. The rate of application is variable according to insect pressure, timing of spray and field scouting. In New York State this product may not be applied within 100 feet (using ground equipment) to 300 feet (using aerial equipment) of coastal marshes or streams that drain into coastal marshes.

Hero EW Insecticide contains the pyrethroids zetacypermethrin and bifenthrin.

Maximum Usage When Applying Zeta-Cypermethrin and Cypermethrin Products to the Same Crop Within the Same Season.

Do not apply more than the maximum seasonal total for either zetacypermethrin or cypermethrin products when used alone; do not apply more than the combined maximum seasonal total for both products as outlined in the table of the label.

#### Limitations, Restrictions, and Exceptions

#### HEAD LETTUCE

##### RESTRICTIONS - Head Lettuce

Refer to the maximum usage tables when applying more than one product containing either zeta-cypermethrin or bifenthrin to this crop.

Do not make applications less than 7 days apart.

##### REMARKS - Head Lettuce

Apply in water as necessary for insect control using a minimum of 15 gallons of finished spray per acre with ground equipment and 5 gallons per acre by air.

When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray.

Thorough coverage is essential to achieve control.

Armyworm fall: Coverage is essential for control of this pest. Under heavy outbreaks tank mixing with another product that is labeled for this pest is recommended for control.

Diamondback Moth, Tobacco Budworm: Pyrethroid resistance is common for this pest. Please consult your local or state agricultural authority to determine if resistance pest populations are in your area. If so refer the resistance management statement in the DIRECTION FOR USE section of the label.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

7 days

Rates

[field\\_rates 0](#)

[field\\_rates 1](#)

•

Restricted Entry Interval

12 hours

Timings

[N.A.](#)