

FLORIDA - ALEXANDERGRASS, ETC.

General Information

GENERAL INFORMATION

Atrazine 90 WSP Herbicide will control most annual Broadleaf and Grass weeds in Corn, Sorghum, Sugarcane and certain other crops as specified on the label. This product may be applied before or after weeds emerge.

In each case where a range of rates is given, the lower rate should be used on Coarse-textured soil and soils low in organic matter and the higher rate should be used on Fine-textured soil and soils high in organic matter. Following many years of continuous use of this product and chemically-related products, biotypes of some of the weeds listed on the label have been reported which cannot be effectively controlled by this and related herbicides. Where this is known or suspected and weeds controlled by this product are expected to be present along with resistant biotypes, we recommend the use of this product in combinations or sequence with other registered herbicides which are not triazines. If only resistant biotypes are expected to be present, use a registered non-triazine herbicide. Consult with your State Agricultural Extension Service for specific recommendations.

Since this product acts mainly through root absorption, its effectiveness depends on moisture to move it into the root zone. Should weeds develop, a shallow cultivation or rotary hoeing will generally result in better weed control.

This product is non-corrosive to equipment and metal surfaces, non-flammable, and has low electrical conductivity. Avoid using this product near adjacent desirable plants or in greenhouses or injury may occur. To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, as crop injury may result.

Note: The Seller does not recommend the use of this product in combination with other herbicides or oils except as specifically described on the label or in literature published by the Seller.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THE LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY OR ILLEGAL RESIDUES.

Do not apply this product through any type of irrigation system.

APPLICATION PROCEDURES

GROUND APPLICATION: Use conventional ground sprayers equipped with nozzles that provide accurate and uniform application. Be certain that nozzles are uniformly spaced and are the same size. Calibrate sprayer before use and recalibrate at the start of each season and when changing carriers. Unless otherwise specified, use a minimum of 10 gallons of spray mixture per acre for all pre-plant incorporated, preplant surface, pre-emergence and post-emergence applications (with or without oil or surfactant) with ground equipment.

Use a pump with capacity to: 1) Maintain 35 to 40 psi at nozzles; 2) Provide sufficient agitation in tank to keep mixture in suspension; and 3) To provide a minimum of 20% bypass at all times.

Use centrifugal pumps which provide propeller shear action for dispersing and mixing this product. The pump should provide a minimum of 10 gallons per minute per 100 gallon tank size circulated through a correctly-positioned sparger tube or jets. Use screens to protect the pump and to prevent nozzles from clogging. Screens placed on the suction side of the pump should be 16-mesh or coarser. Do not place a screen in the recirculation line. Use 50-mesh or coarser screens between the pump and boom and where required, at the nozzles. Check nozzle manufacturer's recommendations.

AERIAL APPLICATION: Use aerial application only where broadcast applications are specified. Apply in a minimum of 1 gallon of water for each pound of this product applied per acre. For post-emergence treatments on Corn and Sorghum, apply recommended rate in a minimum of 2 gallons of water per acre. Avoid applications under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur.

Avoid application to humans or animals. Flagmen and loaders should avoid inhalation of spray mist and prolonged contact with skin.

Limitations, Restrictions, and Exceptions

For control of Alexandergrass, Large crabgrass, Pellitory weed (Artillery weed) and

Spiny amaranth, use one of the following methods at-planting or ratooning:

1. Apply 4.4 pounds of this product per acre [1 pak per 5.6 acres] pre-emergence. Follow with one or two applications, as needed, post-emergence to Sugarcane and weeds at 2.2 pounds of this product per acre [1 pak per 11.3 acres]. Treat before weeds exceed 1.5 inches in height.

2. Apply 1 to 3 times, as needed, at 2.2 pounds of this product per acre [1 pak per 11.3 acres] post-emergence to Sugarcane and weeds. Treat before weeds exceed 1.5 inches in height.

The total pounds of active ingredient applied pre-emergent PLUS post-emergent applications cannot exceed 10 lbs. a.i. (11.1 pounds of this product) [1 pak per 2.2 acres] per acre per crop.

Method

[Broadcast/Foliar Ground](#)

[Directed](#)

[Broadcast/Foliar Ground](#)

[Directed](#)

Rates

[field rates 0](#)

[field rates 1](#)

•

Restricted Entry Interval

12 hours

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[At-Plant](#)

[Ratooning](#)