

FORESTS AND NON-CROPLAND AREAS

General Information

General Information

Pathway herbicide is a ready-to-use product that is effective in cut surface applications for killing unwanted trees and preventing undesirable sprouting of cut trees in forest and other non-crop areas such as fence rows, roadsides and rights-of-way.

General Use Precautions

Not for sale, distribution or use in Nassau and Suffolk Counties in New York State

Use this product only as specified on this label.

Maximum Use Rates: Total use of Pathway must not exceed 4 gallons per acre per annual growing season on rights-of-way and other non-crop areas. No more than 4 gallons per acre may be applied within a period of 2 annual growing seasons on forest sites.

Be sure that use of this product conforms to all applicable regulations. Pathway is highly active as a herbicide, is water-soluble, can move with surface runoff water, and can remain phytotoxic for a year or more if it gets into the soil.

Pathway should not be applied on residential or commercial lawns or near ornamental trees and shrubs. Untreated trees can occasionally be affected by root uptake of herbicide through movement into the topsoil or by excretion of the product from the roots of nearby treated trees. Do not apply Pathway within the root zone of desirable trees unless such injury can be tolerated.

Do not contaminate cropland, water, or irrigation ditches.

Do not contaminate water intended for irrigation or domestic purposes. To avoid injury to crops or other desirable plants, do not treat or allow spray drift or run-off to fall onto banks or bottoms of irrigation ditches, either dry or containing water, or other channels that carry water that may be used for irrigation or domestic purposes. Do not apply to snow or frozen ground.

Do not rotate food or feed crops on treated land if they are not registered for use with picloram until an adequately sensitive bioassay or chemical test shows that no detectable picloram is present in the soil.

Do not apply on or in the vicinity of sensitive crops of desirable broadleaf plants. Small amounts can cause injury to plants.

Do not allow careless application or spray drift. Do not permit any spray or spray drift to contact desirable plants. Use only gravity flow or very low spray pressure.

Limitations, Restrictions, and Exceptions

Application Methods

Tree Injection Treatment

Use 1 milliliter of undiluted Pathway solution through the bark completely around the tree trunk at intervals of 2 to 3 inches between edges of the injector wounds. Make injections near ground level when using the tree injector or 2 to 4 feet above the ground when using a Hypohatchet Injector or similar device. Treatments can be made in any season. Maples should not be treated during the spring sap flow. With some difficult to control species such as dogwood, hickory, sugar maple, bigleaf maple, tanoak, and some firs, application to a continuous cut rather than to spaced cuts may be more uniformly effective. Note: No Worker Protection Standard worker entry restrictions or worker notification requirements apply when this product is directly injected into agricultural plants.

Frill or Girdle Treatment

Make a single hack girdle or "frill" of overlapping ax cuts through the bark completely around the tree as close to the ground as feasible. Spray or paint the injured surface with undiluted Pathway, using enough volume to wet treated areas.

Stump Treatment

Spray or paint the cut surfaces of freshly cut stumps and stubs with undiluted Pathway. The cambium area next to the bark is the most vital area to wet.

Method

[Cut Stump](#)

[Spray](#)

[Tree Injection](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

48 hours

Timings

[Any season](#)