

CORN (FIELD CORN, FIELD CORN GROWN FOR SEED AND POPCORN) - GRAIN AND FODDER

General Information

CORN (FIELD CORN, FIELD CORN GROWN FOR SEED AND POPCORN) - FORAGE

Limitations, Restrictions, and Exceptions

CORN (FIELD CORN, FIELD CORN GROWN FOR SEED AND POPCORN)

Application Timing

Apply when disease first appears and continue on a 10-14 day interval if favorable conditions for disease development persist. Use of shorter spray intervals and higher rates are recommended when disease pressure is severe.

Notes

Absolute Maxx may be applied by ground, air or chemigation.

Absolute Maxx should be applied in a minimum of 10 gallons of spray solution by ground sprayer or in a minimum of 2 gallons per acre by aircraft spray equipment. For optimum disease control, the lowest labeled rate of a spray surfactant may be tank-mixed.

Restrictions: Absolute Maxx may be applied up to 36 days before the harvest of grain and fodder. Do not apply within 21 days of harvest for forage. Do not apply more than two sequential applications of Absolute Maxx. Limit the number of Absolute Maxx or other Group 11-containing fungicide applications to no more than two per acre per crop.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

36 DAYS

-

Restricted Entry Interval

12 HOURS

Timings

[When disease first appears](#)