

COTTON - HIGH RATE SINGLE OR SPLIT APPLICATIONS - FIRST

General Information

GENERAL INFORMATION

HELENA PGR-IV PLUS applications are designed to promote root growth, square retention and boll fill.

Refer to COMPATIBILITY section for tank-mixing with other products. The following table gives the minimum and maximum amounts of diluent to be used to apply the amount of HELENA PGR-IV PLUS per acre specified in the use rate tables.

Applicators should use a dilution rate that achieves thorough coverage of available plant foliage up to the point of runoff.

Limitations, Restrictions, and Exceptions

COTTON

First

Apply lower rate between match-head square and early bloom when planning a sequential application in 18-24 days. Apply higher rate when only a single application is planned.

Restrictions and Limitations

- For use on short-staple and long-staple (Pima) cotton only.
- Do not graze or feed cotton forage to livestock.
- Do not plant another crop within 75 days after last treatment.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

30 days

Rates

[field_rates 0](#)

-

Restricted Entry Interval

12 hours

Timings

[Lower rate: Between match-head square and early bloom.](#)

[Higher rate: Single application.](#)