

CEREAL GRAINS - CORN - BUDWORM, BORER, ARMYWORM ETC.

General Information

GENERAL DIRECTIONS FOR USE

Initial and residual control are contingent upon thorough crop coverage. Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gals./A by air or 10 gals./A by ground unless otherwise specified in the label. When foliage is dense or pest pressure is high (heavier insect or egg pressure, larger larval stages), use of higher application volumes and/or higher use-rates may improve initial and residual control.

For cutworm control, LAMBDA T may be applied before, during, or after planting. For soil incorporated applications, use higher rates for improved control.

Limitations, Restrictions, and Exceptions

CEREAL GRAINS

Corn

- Apply as required by scouting, or locally prescribed corn growth stages, usually at intervals of 7 or more days. Timing and frequency of applications should be based upon insect populations reaching locally-determined economic thresholds or other locally recommended methods.
- Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of target location. When applying by air, apply in a minimum of 2 gals. of water/A.
- For control of adult corn rootworm beetles (*Diabrotica* species) as part of an aerial applied corn rootworm control program use a minimum of 3.84 fl. oz./A (0.03 lb. a.i./A).
- Do not allow livestock to graze in treated areas or harvest treated corn forage as feed for meat or dairy animals within 1 day after last treatment. Do not feed treated corn fodder or silage to meat or dairy animals within 21 days after last treatment.

- Do not apply more than 0.06 lb. a.i. (0.48 pts.) after silk initiation. Do not apply more than 0.03 lb. a.i. (0.24 pts.) after corn has reached the milk stage (yellow kernels with milky fluid).

Target Pest:

Budworm, European Corn Borer, Southwestern Corn Borer, Stalk Borer, Hop Vine Borer: For control before the larva bores into the plant stalk or ear.

Southwestern Corn Borer, Fall Armyworm, Yellow-striped Armyworm²: Use higher rates for large larvae.

Corn Leaf Aphid, Bird Cherry-Oat Aphid, English Grain Aphid: Suppression only.

Budworm: See resistance statement under General Directions for Use.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

21 days

Rates

[field rates 0](#)

[field rates 1](#)

•

Restricted Entry Interval

24 hours

Timings

[At-Plant](#)

[Apply as required by scouting, or locally prescribed corn growth stages, usually at intervals of 7 or more days.](#)