

ORNAMENTALS

General Information

APPLICATION TO ORNAMENTALS

PRO-MATE MERIT 0.2% PLUS TURF FERTILIZER can be used as a broadcast application on ornamentals in commercial and residential landscapes, as well as on landscaped areas of golf courses and other recreational areas. This is a systemic product, with the active ingredient being absorbed by the roots and translocated upward within the plant. Optimum control will be achieved when applications are followed by sufficient mechanical incorporation, irrigation or rainfall to place the active ingredient into the soil.

Applications of PRO-MATE MERIT 0.2% PLUS TURF FERTILIZER cannot exceed a total of 200 lb (0.4 lb of active ingredient) per acre per year.

NOTE: Not for use in commercial greenhouses, or nurseries.

APPLICATION EQUIPMENT

Apply PRO-MATE MERIT 0.2% PLUS TURF FERTILIZER uniformly over the area being treated with typical granular application equipment. Both drop-type and rotary-type spreaders may be used to apply PRO-MATE MERIT 0.2% PLUS TURF FERTILIZER. Avoid the use of spreaders which would apply the material in narrow, concentrated bands. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly and accurately.

SPREADER SETTINGS FOR PRO-MATE MERIT 0.2% PLUS TURF FERTILIZER

The following chart can be used as a guide in establishing an accurate rate of PROMATE MERIT 0.2% PLUS TURF FERTILIZER. The suggested settings should provide an application rate of 3.5 lb per 1000 sq ft, or 150 lb per acre. Application rates will vary dependent upon speed, equipment wear and condition of spreader. Carefully calibrate all equipment used prior to commercial use of this product.

Limitations, Restrictions, and Exceptions

FOR USE ON ORNAMENTALS

Apply as a broadcast treatment. Prior to application, remove black plastic or other weed barrier material that will not permit water permeability.

NOTE: For optimum control, mechanical incorporation, irrigation or rainfall should occur within 24 hours after application to move the active ingredient into soil. Refer to label sections on application equipment and spreader settings.

Method

[Broadcast](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

12 hours

Timings

[Application should be made prior to anticipated pest infestation to achieve optimum levels of control.](#)