

# **CITRUS: CLEMENTINE MANDARIN**

## General Information

### GENERAL USE INSTRUCTIONS

Use only as directed. Read the label thoroughly and understand it before making applications. Keep out of reach of children.

Do not apply this product through any type of irrigation system, unless otherwise permitted on the label.

### Application Instructions

ProGibb LV Plus Plant Growth Regulator (hereafter referred to as ProGibb LV Plus) contains gibberellic acid, which is an extremely potent plant growth regulator; when applying plant growth regulators, deviations from the label directions in the rates, timings, water volumes, or the adoption of untested spray mixes, results in undesirable effects. Always consult the Valent agricultural specialist in your area for the spray regimen best suited to your conditions.

- Do not apply to plants under pest, nutritional, or water stress.
- When a range of rates is indicated, use the concentration and spray volume directed locally by the Valent agricultural specialist.
- For optimum effectiveness, thorough spray coverage must be achieved; all parts of the plant or crop must receive the spray or desired results will not occur. Prepare solution concentrations by mixing the required amount of product with water in a clean, empty spray tank. Dispose of any unused spray material at the end of each day following local, state or federal law.
- For most efficacious results, use water with a pH of 4.0 to 8.5. Use a buffer for water with pH above or below this range.
- ProGibb LV Plus applications made under slow drying conditions (cool to warm temperatures, medium to high relative humidity, and no wind) will increase absorption by the plant, thus optimizing effectiveness. Night time applications are encouraged when day time conditions are not conducive to slow drying conditions.
- Rain fastness: Re-apply ProGibb LV Plus if significant rain occurs within 2 hours of application.
- Compatibility: When considering tank mixing with other products, use the

following compatibility jar test before mixing a whole tank.

- DO NOT apply using ULV application methods. For aerial applications spray volumes must be greater than 2 gallons per acre (10 gallons per acre for tree crops).
- No preharvest interval is required for this product.

#### Compatibility With Other Agricultural Products

Compatibility and performance data for ProGibb LV Plus with other agricultural products are not necessarily available.

Do not tank mix ProGibb LV Plus with other products unless compatibility has been verified. If considering tankmixing ProGibb LV Plus with other products use the following compatibility jar test before mixing a whole tank:

Add water from the same water source to a clear glass or plastic jar. Add the pesticides in correct proportions. Mix thoroughly and let stand for a minimum 15 minutes. Separation, gelling, or generation of heat are all signs of incompatibility.

Even if a mix passes the jar test for compatibility, it is imperative to test it on a designated area to evaluate for phytotoxicity or ineffectiveness.

Always read and follow all label directions and precautions of each product. When using combinations of products the most restrictive of label limitations and precautions must be followed. Do not mix with any pesticide that has a prohibition against tank mixing. For further information consult your Valent agricultural specialist.

#### Directions For Chemigation

Fill the supply tank with the desired amount of water. Then add the amount of ProGibb LV Plus required in order to achieve the final solution rate recommended for the specific crop to be treated. Agitate the mixture of ProGibb LV Plus frequently during the chemigation period to assure a uniform distribution throughout the system. Apply ProGibb LV Plus continuously for the duration of the water application but do not exceed recommended rates and volumes as outlined on the product label.

#### Chemigation Precautions

Apply this product only through the following systems: Overhead sprinklers such as impact, micro-sprinklers, or booms.

Do not apply this product through any other type of irrigation system. Crop injury or lack of effectiveness can result from non-uniform distribution of treated water. If you have any questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. Do not connect an irrigation system(including greenhouse systems) used for pesticide application to a public water system unless the pesticide label prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Prior to application ensure that the chemigation system meets the following requirements: The system must contain a functional check valve, vacuumrelief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

In addition to the above use rates and recommendations, the following precautions must be observed when using this product in any type of irrigation system:

#### Chemigation Systems Connected To Public Water Systems

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days of the year. Chemigation systems connected to public water systems must contain a functional, reduced pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water systems should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where the pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

#### Limitations, Restrictions, and Exceptions

##### CITRUS

For citrus, apply in sprays of sufficient water volumes to ensure thorough fruit wetting. In most cases, this application will cause some drop of older mature leaves;

this drop of older leaves is inconsequential. However, application to trees of low vigor or under stress (pest, nutritional, or water, etc.) have sometimes caused severe leaf and/or fruit drop. Do not apply in white wash sprays in which lime or other caustic material has produced a high pH in the spray tank. Applications of copper fungicides and/or oils within three weeks (before or after) the ProGibb LV Plus application have been known to result in significant leaf drop and fruit drop.

## CLEMENTINE MANDARIN

### OBJECTIVE/BENEFIT

- To increase fruit set and yield.

### APPLICATION TIMING

- Make 1-4 applications from early bloom up to 4 weeks after petal fall. Allow a minimum of 3 days between sprays. Use a dilute spray with sufficient spray volume for adequate coverage of tree canopy.

### NOTE:

- The number of applications depends upon amount of desired fruit set. Generally, more fruit will be set by 2 applications, earlier applications, higher rates, and climatic conditions more favorable to set. Differences in the crop strain have been observed to interact with the above factors to affect the degree of fruit set achieved. Reductions in final fruit size have on occasion occurred as a result of excessive fruit set.

### Method

[Dilute Spray](#)

### Rates

[field\\_rates 0](#)

- 

### Restricted Entry Interval

4 hours

EXCEPTION: If the product is soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

### Timings

From early bloom up to 4 weeks after petal fall