

FRUIT CROPS - BANANAN (NOT FOR USE IN CALIFORNIA) - POSTHARVEST

General Information

GENERAL DIRECTIONS FOR USE

Use only as directed. The label should be read thoroughly and understood before making applications. Keep out of reach of children.

Do not apply this product through any type of irrigation system.

Application Instructions

ProGibb Plus 2X contains gibberellic acid, which is an extremely potent plant growth regulator; when applying plant growth regulators, deviations from the label directions in the rates, timings, water volumes, or the adoption of untested spray mixes, may result in undesirable effects. Always consult the Valent Agricultural Specialist in your area for the spray regimen best suited to your conditions.

- Do not apply to plants under pest, nutritional, or water stress. - When a range of rates is indicated, use the concentration and spray volume recommended locally by the Valent agricultural specialist.
- For optimum effectiveness, thorough spray coverage must be achieved; all parts of the plant or crop must receive the spray or desired results will not occur. Prepare solution concentrations by mixing the required amount of product with water in a clean, empty spray tank. Dispose of any unused spray material at the end of each day following local, state or federal law.
- For best results, the water pH should be around 7.0, and always below 8.5.
- ProGibb applications made under slow drying conditions (cool to warm temperatures, medium to high relative humidity, and no wind) will increase absorption by the plant, thus optimizing effectiveness. Night-time applications are encouraged when day-time conditions are not conducive to slow drying conditions.
- Product persistence: Re-apply ProGibb if significant rain occurs within 2 hours of

application.

- Compatibility: Except when specifically noted, these ProGibb spray guidelines refer to the use of the product alone. The use of surfactants and other additives has been reported to be beneficial. Data concerning the compatibility of ProGibb with other agricultural compounds, except DiPel DF and XenTari DF, are not available. Valent does not assume responsibility for unexpected results due to the tank mixing of ProGibb Plus 2X with other products not recommended on the label.
- DO NOT apply using ULV application methods. For aerial applications spray volumes must be greater than 2 gallons per acre (10 gallons per acre for tree crops).
- No preharvest interval is required for this product.

Limitations, Restrictions, and Exceptions

Post-harvest treatment to extend fruit green life.

The solution can be sprayed or brushed to the crown.

Apply after washing the fruit and before packing. It is permissible to tankmix with other protectants.

NOTE:

- Color development and harvest date may be slightly delayed.
- Use higher rates with heavier crop loads.

Rates

[field rates 0](#)

-

Restricted Entry Interval

4 hours